

A CHARACTERIZATION OF BERGMAN SPACES ON THE UNIT BALL OF \mathbb{C}^n , II

SONGXIAO LI, HASI WULAN, AND KEHE ZHU

ABSTRACT. It was shown in [2] that a holomorphic function f in the unit ball \mathbb{B}_n of \mathbb{C}^n belongs to the weighted Bergman space A_α^p , $p > n + 1 + \alpha$, if and only if the function $|f(z) - f(w)|/|1 - \langle z, w \rangle|$ is in $L^p(\mathbb{B}_n \times \mathbb{B}_n, dv_\beta \times dv_\beta)$, where $\beta = (p + \alpha - n - 1)/2$ and $dv_\beta(z) = (1 - |z|^2)^\beta dv(z)$. In this paper we consider the range $0 < p < n + 1 + \alpha$ and show that in this case $f \in A_\alpha^p$ if and only if the function $|f(z) - f(w)|/|1 - \langle z, w \rangle|$ is in $L^p(\mathbb{B}_n \times \mathbb{B}_n, dv_\alpha \times dv_\alpha)$ if and only if the function $|f(z) - f(w)|/|z - w|$ is in $L^p(\mathbb{B}_n \times \mathbb{B}_n, dv_\alpha \times dv_\alpha)$. We think the revealed difference in the weights for the double integrals between the cases $0 < p < n + 1 + \alpha$ and $p > n + 1 + \alpha$ is somewhat surprising.

1. INTRODUCTION

Let \mathbb{B}_n be the open unit ball in \mathbb{C}^n and dv be the normalized volume measure on \mathbb{B}_n . For any $\alpha > -1$ we introduce the weighted volume measure $dv_\alpha(z) = c_\alpha(1 - |z|^2)^\alpha dv(z)$, where c_α is a positive constant such that $v_\alpha(\mathbb{B}_n) = 1$ again.

For $0 < p < \infty$ and $\alpha > -1$ we use the notation

$$A_\alpha^p = H(\mathbb{B}_n) \cap L^p(\mathbb{B}_n, dv_\alpha)$$

to denote the weighted Bergman spaces on \mathbb{B}_n , where $H(\mathbb{B}_n)$ is the space of all holomorphic functions in \mathbb{B}_n .

It was shown in [2] that for $p > n + 1 + \alpha$, we have $f \in A_\alpha^p$ if and only if

$$\int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \frac{|f(z) - f(w)|^p}{|1 - \langle z, w \rangle|^p} dv_\beta(z) dv_\beta(w) < \infty,$$

where $\beta = (p + \alpha - n - 1)/2$. An example in [3] shows that this is no longer true in general when $p = n + 1 + \alpha$.

The purpose of this article is to examine the remaining case $0 < p < n + 1 + \alpha$. Our main result is the following.

2000 *Mathematics Subject Classification.* 32A36.

Key words and phrases. Bergman spaces, unit ball, volume measure.

All authors were partially supported by the China National Natural Science Foundation.

Zhu was partially supported by the US National Science Foundation.

Theorem A. *Suppose $\alpha > -1$, $0 < p < \alpha + n + 1$, and f is holomorphic in \mathbb{B}_n . Then the following conditions are equivalent.*

- (a) $f \in A_\alpha^p(\mathbb{B}_n)$.
 (b) *The double integral*

$$\int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \frac{|f(z) - f(w)|^p}{|z - w|^p} dv_\alpha(z) dv_\alpha(w)$$

converges.

- (c) *The double integral*

$$\int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \frac{|f(z) - f(w)|^p}{|1 - \langle z, w \rangle|^p} dv_\alpha(z) dv_\alpha(w)$$

converges.

An interesting special case is obtained when $n = 1$, $\alpha > 0$, and $p = 2$. In this case, the unit ball \mathbb{B}_n becomes the unit disk \mathbb{D} , and membership in the Bergman spaces $A_\alpha^2(\mathbb{D})$ and $A_\alpha^2(\mathbb{D}^2)$ can be described by Taylor coefficients. As a consequence, we see that for $\alpha > 0$,

$$\sum_{i=1}^k \frac{1}{i^{\alpha+1}(k+1-i)^{\alpha+1}} \sim \frac{1}{k^{\alpha+1}}$$

as $k \rightarrow \infty$. This does not seem to be obvious. Once again, this is not true when $\alpha = 0$.

2. PRELIMINARY ESTIMATES

We begin with an identity concerning Möbius maps on the unit ball and derive some useful inequalities. This will play an important role in the proof of Theorem A.

Lemma 1. *For all z and w in \mathbb{B}_n we have*

$$(1) \quad |z - \varphi_z(w)|^2 = \frac{(1 - |z|^2)(|w|^2 - |\langle z, w \rangle|^2)}{|1 - \langle z, w \rangle|^2}.$$

Consequently,

$$(2) \quad |z - \varphi_z(w)| \geq \frac{|w|(1 - |z|^2)}{|1 - \langle z, w \rangle|},$$

and

$$(3) \quad |z - \varphi_z(w)|^2 \leq \frac{2(1 - |z|^2)}{|1 - \langle z, w \rangle|}.$$

Proof. To prove the identity in (1), we use Lemmas 1.2 and 1.3 in [5] along with the fact that $z = \varphi_z(0)$. Let $S = |z - \varphi_z(w)|^2$. Then

$$\begin{aligned}
S &= |z|^2 + |\varphi_z(w)|^2 - \langle z, \varphi_z(w) \rangle - \langle \varphi_z(w), z \rangle \\
&= |z|^2 + |\varphi_z(w)|^2 - \left(1 - \frac{1 - |z|^2}{1 - \langle z, w \rangle}\right) - \left(1 - \frac{1 - |z|^2}{1 - \langle w, z \rangle}\right) \\
&= -(1 - |z|^2) - (1 - |\varphi_z(w)|^2) + \frac{1 - |z|^2}{1 - \langle z, w \rangle} + \frac{1 - |z|^2}{1 - \langle w, z \rangle} \\
&= \frac{1 - |z|^2}{1 - \langle z, w \rangle} + \frac{1 - |z|^2}{1 - \langle w, z \rangle} - (1 - |z|^2) - \frac{(1 - |z|^2)(1 - |w|^2)}{|1 - \langle z, w \rangle|^2} \\
&= (1 - |z|^2) \left[\frac{1}{|1 - \langle z, w \rangle|^2} - \left|1 - \frac{1}{1 - \langle z, w \rangle}\right|^2 - \frac{1 - |w|^2}{|1 - \langle z, w \rangle|^2} \right] \\
&= \frac{(1 - |z|^2)(|w|^2 - |\langle z, w \rangle|^2)}{|1 - \langle z, w \rangle|^2}.
\end{aligned}$$

This proves the identity in (1). The inequality in (2) follows from (1) and the fact that $|\langle z, w \rangle| \leq |z||w|$. The inequality in (3) follows from (1) and the following estimates:

$$\begin{aligned}
|w|^2 - |\langle z, w \rangle|^2 &\leq 1 - |\langle z, w \rangle|^2 \\
&= (1 + |\langle z, w \rangle|)(1 - |\langle z, w \rangle|) \\
&\leq 2|1 - \langle z, w \rangle|.
\end{aligned}$$

This completes the proof of the lemma. \square

Note that Lemma 1 above appears as Exercise 3.18 in [5]. We believe it never appeared elsewhere before, so a full proof is given here.

Lemma 2. *There exists a positive constant C , independent of f , such that*

$$\int_{\mathbb{B}_n} \frac{|f(w)|^p}{|w|^p} dv_\alpha(w) \leq C \int_{\mathbb{B}_n} |f(w)|^p dv_\alpha(w)$$

for all $f \in H(\mathbb{B}_n)$ with $f(0) = 0$.

Proof. This is well known. See Lemma 4.26 of [4] in the one-dimensional case. The higher dimensional case is proved in a similar way. \square

Lemma 3. *For any $r > 0$ and $z \in \mathbb{B}_n$ let $D(z, r)$ be the Bergman metric ball at z .*

(a) *There exists a positive constant C (dependent on r , p , and α , but not on f and z) such that*

$$|f(z)|^p \leq \frac{C}{(1 - |z|^2)^{n+1+\alpha}} \int_{D(z, r)} |f(w)|^p dv_\alpha(w)$$

for all $f \in H(\mathbb{B}_n)$ and $z \in \mathbb{B}_n$.

(b) There exists a positive constant C (dependent on r but not on z and w) such that

$$C^{-1}(1 - |w|^2) \leq |1 - \langle z, w \rangle| \leq C(1 - |z|^2)$$

for all $z \in \mathbb{B}_n$ and $w \in D(z, r)$.

Proof. This is also well known. See Lemma 2.24 of [5] for example. \square

Lemma 4. Suppose $\alpha > -1$ and $t < 0$. Then there exists a positive constant C such that

$$\int_{\mathbb{B}_n} \frac{dv_\alpha(w)}{|1 - \langle z, w \rangle|^{n+1+\alpha+t}} \leq C$$

for all $z \in \mathbb{B}_n$.

Proof. This is part of Theorem 1.12 in [5]. \square

3. MAIN RESULT

We now proceed to prove Theorem A, the main result of the paper. This will be accomplished by the following three lemmas.

Lemma 5. There exists a positive constant C such that $I(f) \leq CJ(f)$ for all $f \in H(\mathbb{B}_n)$, where

$$I(f) = \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \frac{|f(z) - f(w)|^p}{|z - w|^p} dv_\alpha(z) dv_\alpha(w),$$

and

$$J(f) = \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \frac{|f(z) - f(w)|^p}{|1 - \langle z, w \rangle|^p} dv_\alpha(z) dv_\alpha(w).$$

Proof. By Fubini's theorem and a change of variables (see Proposition 1.13 of [5]),

$$I(f) = \int_{\mathbb{B}_n} dv_\alpha(z) \int_{\mathbb{B}_n} \frac{|f(z) - f \circ \varphi_z(w)|^p}{|z - \varphi_z(w)|^p} \frac{(1 - |z|^2)^{n+1+\alpha}}{|1 - \langle z, w \rangle|^{2(n+1+\alpha)}} dv_\alpha(w).$$

By (2), there exists a positive constant C such that $I(f)$ is less than or equal to

$$C \int_{\mathbb{B}_n} (1 - |z|^2)^{n+1+\alpha-p} dv_\alpha(z) \int_{\mathbb{B}_n} \frac{|f(z) - f \circ \varphi_z(w)|^p}{|w|^p |1 - \langle z, w \rangle|^{2(n+1+\alpha)-p}} dv_\alpha(w).$$

By Lemma 2, there exists another constant $C > 0$ such that $I(f)$ is less than or equal to

$$C \int_{\mathbb{B}_n} (1 - |z|^2)^{n+1+\alpha-p} dv_\alpha(z) \int_{\mathbb{B}_n} \frac{|f(z) - f \circ \varphi_z(w)|^p}{|1 - \langle z, w \rangle|^{2(n+1+\alpha)-p}} dv_\alpha(w).$$

Another change of variables then gives

$$I(f) \leq C \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \frac{|f(z) - f(w)|^p}{|1 - \langle z, w \rangle|^p} dv_\alpha(z) dv_\alpha(w) = CJ(f).$$

This proves the lemma. \square

Lemma 6. *There exists a positive constant C such that*

$$\int_{\mathbb{B}_n} |f(z) - f(0)|^p dv_\alpha(z) \leq C \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \frac{|f(z) - f(w)|^p}{|z - w|^p} dv_\alpha(z) dv_\alpha(w)$$

for all $f \in H(\mathbb{B}_n)$.

Proof. We use $I(f)$ again to denote the integral on the right hand side of the inequality to be proved. And again we make a change of variables to obtain

$$I(f) = \int_{\mathbb{B}_n} dv_\alpha(z) \int_{\mathbb{B}_n} \frac{|f(z) - f \circ \varphi_z(w)|^p}{|z - \varphi_z(w)|^p} \frac{(1 - |z|^2)^{n+1+\alpha}}{|1 - \langle z, w \rangle|^{2(n+1+\alpha)}} dv_\alpha(w).$$

By the inequality in (3) of Lemma 1, there exists a positive constant C such that $I(f)$ is greater than or equal to

$$C \int_{\mathbb{B}_n} dv_\alpha(z) \int_{\mathbb{B}_n} \frac{|f(z) - f \circ \varphi_z(w)|^p}{\frac{(1 - |z|^2)^{p/2}}{|1 - \langle z, w \rangle|^{p/2}}} \frac{(1 - |z|^2)^{n+1+\alpha}}{|1 - \langle z, w \rangle|^{2(n+1+\alpha)}} dv_\alpha(w).$$

If we let $D(z, r)$ denote the Bergman metric ball, then $I(f)$ is greater than or equal to

$$C \int_{\mathbb{B}_n} dv_\alpha(z) \int_{D(z, r)} |f(z) - f \circ \varphi_z(w)|^p \frac{(1 - |z|^2)^{n+1+\alpha-p/2}}{|1 - \langle z, w \rangle|^{2(n+1+\alpha)-p/2}} dv_\alpha(w).$$

This together with part (a) of Lemma 3 shows that there is another positive constant C , independent of f and z , such that

$$I(f) \geq C \int_{\mathbb{B}_n} dv_\alpha(z) \int_{D(z, r)} \frac{|f(z) - f \circ \varphi_z(w)|^p}{(1 - |z|^2)^{n+1+\alpha}} dv_\alpha(w).$$

Combining this with part (b) of Lemma 3, we obtain yet another positive constant C such that

$$I(f) \geq C \int_{\mathbb{B}_n} |f(z) - f \circ \varphi_z(z)|^p dv_\alpha(z),$$

or

$$I \geq C \int_{\mathbb{B}_n} |f(z) - f(0)|^p dv_\alpha(z).$$

This proves the lemma. \square

Lemma 7. *If $0 < p < n + 1 + \alpha$ and $f \in A_\alpha^p(\mathbb{B}_n)$, then the double integral*

$$\int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \frac{|f(z) - f(w)|^p}{|z - w|^p} dv_\alpha(z) dv_\alpha(w)$$

converges.

Proof. Once again we use $I(f)$ to denote the double integral above. By the remark at the end of [3], there exists a positive continuous function $g \in L^p(\mathbb{B}_n, dv_\alpha)$ such that

$$|f(z) - f(w)| \leq d(z, w)(g(z) + g(w)), \quad z, w \in \mathbb{B}_n,$$

where

$$d(z, w) = \frac{|z - w|}{|1 - \langle z, w \rangle|}.$$

It follows that

$$I(f) \leq \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \left(\frac{g(z) + g(w)}{|1 - \langle z, w \rangle|} \right)^p dv_\alpha(z) dv_\alpha(w).$$

Choose a positive constant C such that

$$(x + y)^p \leq C(x^p + y^p)$$

for all positive values x and y . Then

$$I(f) \leq C \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \frac{g(z)^p + g(w)^p}{|1 - \langle z, w \rangle|^p} dv_\alpha(z) dv_\alpha(w).$$

This along with Fubini's theorem gives

$$I(f) \leq 2C \int_{\mathbb{B}_n} g(z)^p dv_\alpha(z) \int_{\mathbb{B}_n} \frac{dv_\alpha(w)}{|1 - \langle z, w \rangle|^p}.$$

Since $p < n + 1 + \alpha$, it follows from Lemma 4 that there is another positive constant C such that

$$I(f) \leq C \int_{\mathbb{B}_n} g(z)^p dv_\alpha(z) < \infty.$$

This proves the lemma. \square

Note that Lemma 7 above was obtained in [3] in the one-dimensional case. Combining Lemmas 5-7, we have completed the proof of Theorem A. We can slightly strengthen Theorem A as follows.

Corollary 8. *Fix $\alpha > -1$ and $p > 0$. For any holomorphic function f in \mathbb{B}_n let*

$$\|f\|_1^p = \int_{\mathbb{B}_n} |f(z)|^p dv_\alpha(z),$$

and

$$\|f\|_2^p = |f(0)|^p + \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \frac{|f(z) - f(w)|^p}{|1 - \langle z, w \rangle|^p} dv_\alpha(z) dv_\alpha(w),$$

and

$$\|f\|_3^p = |f(0)|^p + \int_{\mathbb{B}_n} \int_{\mathbb{B}_n} \frac{|f(z) - f(w)|^p}{|z - w|^p} dv_\alpha(z) dv_\alpha(w).$$

Then there exists a positive constant C such that

$$C^{-1}\|f\|_k \leq \|f\|_l \leq C\|f\|_k, \quad 1 \leq l \leq 3, \quad 1 \leq k \leq 3,$$

for all $f \in H(\mathbb{B}_n)$.

Proof. This follows from Theorem A and the open mapping theorem. \square

REFERENCES

- [1] F. Holland and D. Walsh, Criteria for membership of Bloch space and its subspace, BMOA, *Math. Ann.* **273** (1986), 317-335.
- [2] S. Li, H. Wulan, R. Zhao, and K. Zhu, A characterization of Bergman spaces on the unit ball of \mathbb{C}^n , to appear in *Glasgow Math. J.*
- [3] H. Wulan and K. Zhu, Lipschitz type characterizations for Bergman spaces, to appear in *Canadian Math. Bull.*
- [4] K. Zhu, *Operator Theory in Function Spaces*, second edition, American Mathematical Society, Providence, Rhode Island, 2007.
- [5] K. Zhu, *Spaces of Holomorphic Functions in the Unit Ball*, Springer, New York, 2005.

LI: DEPARTMENT OF MATHEMATICS, JIAYING UNIVERSITY, MEIZHOU, GUANGDONG 514015, CHINA

E-mail address: jyulsx@163.com

WULAN: DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU, GUANGDONG 515063, CHINA

E-mail address: wulan@stu.edu.cn

ZHU: DEPARTMENT OF MATHEMATICS AND STATISTICS, STATE UNIVERSITY OF NEW YORK, ALBANY, NY 12222, USA

E-mail address: kzhu@math.albany.edu