

COMBINATORIAL REPRESENTATION THEORY OF LIE ALGEBRAS. RICHARD STANLEY'S WORK AND THE WAY IT WAS CONTINUED

CRISTIAN LENART

Dedicated to Richard Stanley on the occasion of his seventieth birthday.

Representation theory is a fundamental tool for studying group symmetry – geometric, analytic, or algebraic – by means of linear algebra, which has important applications to other areas of mathematics and mathematical physics. One very successful trend in this field in recent decades involves using combinatorial objects to model the representations, which allows one to apply combinatorial methods for studying them, e.g., for concrete computations. This trend led to the emergence of combinatorial representation theory, which has now become a thriving area.

Richard Stanley played a crucial role, through his work and his students, in the development of this new area. In the early stages, he has the merit to have pointed out to combinatorialists, in [34, 36], the potential that representation theory has for applications of combinatorial methods. Throughout his distinguished career, he wrote significant articles which touch upon various combinatorial aspects related to representation theory (of Lie algebras, the symmetric group, etc.). I will describe some of Richard's contributions involving Lie algebras, as well as recent developments inspired by them (including some open problems), which attest the lasting impact of his work.

Acknowledgement. Meeting Richard was a defining moment of my life and career, as well as of my evolution since then. Therefore, I always said that I consider him my mentor. I admire him both as a person and as a scientist, while his vast work in combinatorics and on its many relationships with other areas of mathematics is a continuous inspiration for me.

1. INTRODUCTION

The story starts with the well-known fact that the *Schur symmetric polynomials* $s_\lambda(x_1, \dots, x_n)$, indexed by partitions $\lambda = (\lambda_1 \geq \dots \geq \lambda_n \geq 0)$, are the irreducible polynomial characters of $GL_n = GL_n(\mathbb{C})$. (A representation $X \in GL_n \mapsto \psi(X) \in GL(V)$ of GL_n on a vector space V is polynomial, resp. rational, if the entries of $\psi(X)$ are polynomial, resp. rational functions of the entries of X .) We have the immediate connection to combinatorics, due to the well-known formula:

$$(1) \quad s_\lambda(x_1, \dots, x_n) = \sum_{T \in SSYT_{[n]}(\lambda)} x^T.$$

Here $SSYT_{[n]}(\lambda)$ denotes the set of *semistandard Young tableaux* of shape λ filled with integers in $[n] := \{1, \dots, n\}$, that is, fillings of the Young diagram λ with weakly increasing rows (from left to right) and strictly increasing columns (from top to bottom), where we use the English convention; the monomial x^T is given by $x_1^{c_1} \dots x_n^{c_n}$, where c_i is the number of i 's in T .

The very rich combinatorics of symmetric polynomials (or symmetric functions, if infinitely many variables are used) is beautifully described by Richard in [43][Chapter 7], to which we refer for more information on the topics below. Many questions in the representation theory of GL_n , or the Lie algebra $\mathfrak{sl}_n = \mathfrak{sl}_n(\mathbb{C})$, are answered in combinatorial terms. A typical example is the important problem of decomposing tensor products of irreducible representations; this problem is reduced to

C.L. was partially supported by the NSF grant DMS-1101264, and gratefully acknowledges the hospitality and support of the Max-Planck-Institut für Mathematik in Bonn, where this article was written.

describing the structure constants (or Littlewood-Richardson coefficients) for the multiplication of Schur functions, and its solution is given by the celebrated *Littlewood-Richardson rule*.

2. THE ROBINSON-SCHENSTED-KNUTH CORRESPONDENCE

2.1. The classical correspondence. The *Robinson-Schensted-Knuth (RSK) correspondence* is a well-known bijection between non-negative integer matrices with finite support and pairs of SSYT of the same shape. It has an important role in the theory of symmetric functions; for instance, it bijectively proves the *Cauchy identity* for Schur functions, cf. (2).

Let us briefly explain the RSK correspondence. Given a non-negative integer matrix (a_{ij}) with finite support, we first associate with it (in a bijective way) a biword, that is, a pair of words $(\mathbf{i} = i_1 \dots i_r, \mathbf{j} = j_1 \dots j_r)$, viewed up to permutation of biletters (i_k, j_k) . The bijection is given by considering a_{ij} biletters (i, j) for each $a_{ij} > 0$. Unless otherwise specified, we think of a biword as its lexicographic representative, that is, we order the biletters such that $i_k < i_{k+1}$, or $i_k = i_{k+1}$ and $j_k \leq j_{k+1}$, for all k . We then map (\mathbf{i}, \mathbf{j}) to a pair (P, Q) of SSYT of the same shape by the *Schensted insertion* procedure applied to the word \mathbf{j} (which gives P), run in parallel with a “recording” procedure applied to the word \mathbf{i} (which gives Q). It is well-known that the transpose of the matrix (a_{ij}) , i.e., the biword (\mathbf{j}, \mathbf{i}) , is mapped to (Q, P) .

2.2. A skew analogue. In Richard’s joint paper with B. Sagan [32], a generalization of the RSK correspondence to skew tableaux is constructed.

Given a skew Young diagram λ/μ , for $\mu \subset \lambda$, we denote its semistandard fillings by $SSYT(\lambda/\mu)$, and define the skew Schur function $s_{\lambda/\mu}(x)$ in infinitely many variables by the analogue of (1).

Theorem 2.1. [32][Theorem 6.1] *Let α, β be fixed partitions. There is a bijection $((\mathbf{i}, \mathbf{j}), T, U) \leftrightarrow (P, Q)$ between (lexicographic) biwords (\mathbf{i}, \mathbf{j}) with $T \in SSYT(\alpha/\mu)$, $U \in SSYT(\beta/\mu)$, and $P \in SSYT(\lambda/\beta)$, $Q \in SSYT(\lambda/\alpha)$ such that $\mathbf{i} \cup T = P$ and $\mathbf{j} \cup U = Q$, as unions of the corresponding multisets of entries (defined by adding the number of repetitions of an element).*

The bijection in Theorem 2.1 is based on two types of insertion procedures: an external and an internal insertion, both of which are based on the usual Schensted insertion. The skew RSK correspondence bijectively proves the following generalization of the Cauchy identity (the classical case corresponds to $\alpha = \beta$ being the empty partition):

$$(2) \quad \sum_{\lambda} s_{\lambda/\beta}(x) s_{\lambda/\alpha}(y) = \sum_{\mu} s_{\alpha/\mu}(x) s_{\beta/\mu}(y) \prod_{i,j} (1 - x_i y_j)^{-1}.$$

Other results related to the skew RSK correspondence are derived in [32]: transposing the matrix exchanges the two output tableaux (like in the classical RSK), various identities for $s_{\lambda/\mu}(x)$ and the number of skew standard Young tableaux $f^{\lambda/\mu}$ (a SSYT is standard if its entries are 1 through $n = |\lambda/\mu|$), a skew dual RSK correspondence (corresponding to 0, 1-matrices and output tableaux of conjugate shapes), and an RSK correspondence for so-called skew *shifted tableaux*.

2.3. A connection to representation theory. We will give a representation-theoretic interpretation of the RSK correspondence, in terms of *Kashiwara’s crystals* [15].

Crystals are colored directed graphs encoding representations V of a quantum group $U_q(\mathfrak{g})$, i.e., a certain q -deformation of the enveloping algebra of a Lie algebra \mathfrak{g} , in the limit $q \rightarrow 0$. The vertices of the crystal correspond to the elements b of a so-called *crystal basis* B of V , and the edges give the action on B of the *Kashiwara* (or *crystal*) operators \tilde{f}_i (i.e., certain analogues of the Chevalley generators f_i) in the limit $q \rightarrow 0$. More precisely, whenever B exists, \tilde{f}_i permutes its elements or sends them to 0, so $\tilde{f}_i : B \rightarrow B \cup \{0\}$, and $\tilde{f}_i(b) = b'$ is represented by an i -colored edge $b \xrightarrow{i} b'$.

For instance, the r th tensor power of the vector representation of the Lie algebra \mathfrak{gl}_n has a crystal basis indexed by words of length r in the alphabet $[n]$. By taking the direct limit as $n \rightarrow \infty$, we can

define the $\mathfrak{gl}_{\geq 0}$ -crystal structure on words in the alphabet $\mathbb{Z}_{>0}$. In fact, there is a nice construction of the map f_i on words (where $i \in \mathbb{Z}_{>0}$), based on the so-called *signature rule*. Namely, given a word w , form a binary \pm -word by recording each i (resp. $i+1$) in w as a $+$ (resp. $-$), from left to right. Then cancel adjacent pairs $-+$ as long as possible, to obtain the i -signature of w , which has the form $+\dots+ -\dots-$. If this contains no $+$, then $f_i(w) := \mathbf{0}$; otherwise, $f_i(w)$ is defined by changing to $i+1$ the i in w corresponding to the rightmost $+$ in the i -signature. For instance, $f_2(34321222342) = 3432122\mathbf{3}342$.

It is useful to identify a SSYT T with its word $w(T)$, by reading the entries of T from left to right in each row, with the rows considered from bottom to top. It is well-known that the class of (words of) SSYT is closed under the crystal operators, so it forms a $\mathfrak{gl}_{>0}$ -subcrystal.

We define (commuting) left and right $\mathfrak{gl}_{>0}$ -crystal structures on pairs (P, Q) of SSYT by letting the crystal operators act on P , resp. Q . We can also define a right $\mathfrak{gl}_{>0}$ -crystal structure on biwords (\mathbf{i}, \mathbf{j}) by ordering the biword lexicographically, as above, which gives $(\mathbf{i}_R, \mathbf{j}_R)$, and by letting the crystal operators act on \mathbf{j}_R . Similarly, by exchanging the roles of \mathbf{i} and \mathbf{j} in the definition of the lexicographic order, which gives $(\mathbf{i}_L, \mathbf{j}_L)$, we can define a left $\mathfrak{gl}_{>0}$ -crystal structure on biwords.

Theorem 2.2. [21] *The RSK bijection identifies the left (resp. right) crystal structure on biwords with the right (resp. left) crystal structure on pairs of SSYT. Namely, assuming $(\mathbf{i}, \mathbf{j}) \leftrightarrow (P, Q)$, we have $f_i(\mathbf{j}_R) \neq \mathbf{0}$ if and only if $f_i(P) \neq \mathbf{0}$, in which case*

$$(\mathbf{i}_R, f_i(\mathbf{j}_R)) \leftrightarrow (f_i(P), Q).$$

Similarly, $f_i(\mathbf{i}_L) \neq \mathbf{0}$ if and only if $f_i(Q) \neq \mathbf{0}$, in which case

$$(f_i(\mathbf{i}_L), \mathbf{j}_L) \leftrightarrow (P, f_i(Q)).$$

More generally, one can consider the Lie algebra \mathfrak{gl}_∞ of matrices of finite support with rows and columns indexed by \mathbb{Z}^* , which has $\mathfrak{gl}_{>0}$ as a subalgebra, together with the obvious isomorphic copy $\mathfrak{gl}_{<0}$. In this context, Kwon states Theorem 2.2 in [18] as an isomorphism of $(\mathfrak{gl}_{<0}, \mathfrak{gl}_{>0})$ -crystals; here the alphabet corresponding to $\mathfrak{gl}_{<0}$, which indexes the corresponding crystal operators, is taken to be $\mathbb{Z}_{<0}$. Moreover, an f_0 crystal operator is defined both for biwords and pairs of SSYT, giving \mathfrak{gl}_∞ -crystal structures. Then Theorem 2.2 is strengthened in [18] by stating that the RSK correspondence is a \mathfrak{gl}_∞ -crystal isomorphism. Note, however, that not the original RSK correspondence is used here, but its seventh variation (out of eight) in [9][Appendix A.4.2, Theorem 1].

The RSK bijection is then interpreted in [18] as an isomorphism between the above \mathfrak{gl}_∞ -crystal of matrices and the crystal of a generalized Verma module for \mathfrak{gl}_∞ . Finally, a refinement of this isomorphism is given; a special case of it is precisely the Sagan-Stanley skew RSK correspondence.

A representation-theoretic interpretation of (2), also based on crystals, is given in [19]. Furthermore, an application of the skew dual RSK correspondence in [32] to the construction of crystal bases of certain modules for the quantum superalgebra $U_q(\mathfrak{gl}(m|n))$ is given in [20].

3. \mathfrak{sl}_2 -REPRESENTATIONS AND RELATED POSETS

3.1. Combinatorics from \mathfrak{sl}_2 -representations. The representations of the Lie algebra $\mathfrak{sl}_2 = \mathfrak{sl}_2(\mathbb{C})$ of complex 2×2 matrices of trace 0 are a powerful tool for proving various combinatorial properties of certain posets. They have been used by Richard in [34, 36, 40].

We start by observing that the character of an \mathfrak{sl}_2 -representation on an n -dimensional vector space V , that is, a Lie algebra homomorphism $\psi : \mathfrak{sl}_2 \rightarrow \mathfrak{gl}(V)$, can be expressed as

$$ch(\psi) = \sum_{i=1}^n q^{e_i};$$

here e_i are the eigenvalues of the $n \times n$ matrix $\psi(H)$, with $H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. As it is well-known, the irreducible representations of \mathfrak{sl}_2 are labeled by $k \geq 0$, so we denote them by ψ_k ; we also have

$$ch(\psi_k) = q^{-k} + q^{-k+2} + q^{-k+4} + \dots + q^k.$$

This basic fact implies the following main tool for combinatorics.

Theorem 3.1. [40][Theorem 15] *Let ψ be an \mathfrak{sl}_2 -representation with character $ch(\psi) = \sum_i b_i q^i$. Then the following two sequences are symmetric and unimodal:*

$$\dots, b_{-4}, b_{-2}, b_0, b_2, b_4, \dots \quad \text{and} \quad \dots, b_{-3}, b_{-1}, b_1, b_3, \dots$$

Recall that a sequence a_0, a_1, \dots, a_d is symmetric if $a_i = a_{d-i}$ for all i , and unimodal if, for some i , we have $a_0 \leq a_1 \leq \dots \leq a_i \geq a_{i+1} \geq \dots \geq a_d$. Several applications of Theorem 3.1 were given in [34, 35, 36, 40]. Typical ones involve ranked posets and their rank cardinalities. Interesting examples come from topology, and are intimately related to the *hard Lefschetz theorem*; here the \mathfrak{sl}_2 -representation is on the cohomology or intersection cohomology of an irreducible complex projective variety. Among these examples are the ones involving parabolic quotients of Weyl groups and the generalized lower bound conjecture for simplicial polytopes. These results are surveyed elsewhere in this volume, so we will not address them here.

Another interesting property of a ranked poset which can be proved via \mathfrak{sl}_2 -representations is the *strong Sperner property*, see [29, 35]. This says that the largest subset of the poset containing no $k + 1$ -element chain has the same size as the union of the k largest ranks, for each k (the classical Sperner property corresponds to $k = 1$). As this property is proved in conjunction with the properties discussed above, the following concept is useful: a ranked poset is called *Peck* if it is strongly Sperner and its rank sequence is symmetric and unimodal.

Similar results (unimodality, Sperner property) were derived by Richard in [38] by using the superalgebra analogue of \mathfrak{sl}_2 , namely the orthosymplectic superalgebra $\mathfrak{osp}(1, 2)$.

3.2. Differential posets. As mentioned above, many interesting \mathfrak{sl}_2 -representations were constructed based on posets. This fact led to the concept of an \mathfrak{sl}_2 -poset. To define it, we need some notation. Given a poset P and a field K , let KP denote the K -vector space with basis P . The covering relation in P is denoted by $<$. If P is ranked, let P_i be the set of elements of rank i .

Definition 3.2. [29], [42][Section 4] *A finite ranked poset P of rank n is called an \mathfrak{sl}_2 -poset if there exist operators $X, Y : KP \rightarrow KP$ satisfying*

$$(3) \quad Xx = \sum_{y: y \succ x} c_{xy} y \quad \text{for } x \in P \text{ (where } c_{xy} \in K), \quad Yx \in KP_{i-1} \quad \text{for } x \in P_i,$$

as well as the following commutation relation in every rank i :

$$(4) \quad (XY - YX)_i = (2i - n) Id_i.$$

The operator X is called an order-raising operator, while Y is called a lowering operator. Together they define an \mathfrak{sl}_2 -representation in the usual way, namely $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \mapsto X$ and $\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \mapsto Y$. Using methods similar to those in [42], Proctor proved the following theorem, cf. Section 3.1.

Theorem 3.3. [29][Theorem 1], [42] *A ranked poset P is Peck if and only if it is an \mathfrak{sl}_2 -poset.*

An \mathfrak{sl}_2 -poset is, in fact, one of the variations in [42] of so-called *differential posets*. The latter were defined by Richard in the earlier paper [39], as a generalization of Young's lattice of partitions (under inclusion).

Definition 3.4. [39] A poset P is called r -differential (for some fixed positive integer r) if the operators $U, D : KP \rightarrow KP$ given by

$$(5) \quad Ux := \sum_{y : y > x} y, \quad Dx := \sum_{y : y < x} y$$

satisfy the condition

$$(6) \quad DU - UD = r \text{Id}.$$

Note the similarity of (3) and (4) with (5) and (6), respectively. Many remarkable algebraic and combinatorial properties of differential posets and of their variations are derived by Richard in [39, 42]. Other researchers continued this study. For instance, Fomin [8] proves a generalization of the RSK correspondence, cf. Section 2, to differential posets (in fact, to his slightly more general class of *dual graded graphs*). His related concept of *growth diagrams* (see also [43][Appendix 1]) has proved to be very useful in representation theory in relation to crystals, see [22, 45].

Degree 3 relations between the down and up operators on posets defined in (5) were considered in the literature. Benkart and Roby [1] study the infinite-dimensional associative algebras generated by such operators, which they call *down-up algebras*. They show that these algebras exhibit many of the important features of the universal enveloping algebra of \mathfrak{sl}_2 , including a Poincaré-Birkhoff-Witt type basis and a well-behaved representation theory.

3.3. Representation diagrams. We will now discuss the way in which several \mathfrak{sl}_2 -posets can be pieced together, in order to describe a representation of any semisimple Lie algebra \mathfrak{g} , with Chevalley generators $\{X_i, Y_i, H_i\}_{i \in I}$. These ideas are due to R. Proctor and his student, R. Donnelly, and we refer to [5] for more details.

We start with a finite ranked poset P , viewed as a directed graph with edges $x \rightarrow y$ for each covering relation $x < y$. These edges will be colored by the set I above, so we denote them by $x \xrightarrow{i} y$ for some $i \in I$. The connected components of the subgraph with edges colored i are called *i -components*. Beside a given color, each edge $x \rightarrow y$ is labeled with two complex coefficients, which are not both 0, and which are denoted by c_{yx} and d_{xy} .

Based on the above structure, we define operators X_i and Y_i on $\mathbb{C}P$ for i in I , as follows:

$$(7) \quad X_i x := \sum_{y : x \xrightarrow{i} y} c_{yx} y, \quad Y_i y := \sum_{x : x \xrightarrow{i} y} d_{xy} x.$$

For each vertex x of P , we also define a set of integers $\{m_i(x)\}_{i \in I}$ by $m_i(x) := 2\rho_i(x) - l_i(x)$, where $l_i(x)$ is the rank of the i -component containing x , and $\rho_i(x)$ is the rank of x within that component.

The goal is to define a representation of \mathfrak{g} on $\mathbb{C}P$ by letting the Chevalley generators X_i and Y_i act as in (7), and by setting

$$(8) \quad H_i x := m_i(x) x.$$

Note the similarity of (7) and (8) with (3) and (4), respectively (recall that $H_i = [X_i, Y_i]$). We will give necessary and sufficient local conditions on the edge labels, which are translations of the *Serre relations* for the Chevalley generators of \mathfrak{g} .

Let $\{\omega_i\}_{i \in I}$ and $\{\alpha_i\}_{i \in I}$ denote the fundamental weights and simple roots of the root system corresponding to \mathfrak{g} , respectively. We assign a weight to each vertex of P by $\text{wt}(x) := \sum_{i \in I} m_i(x)\omega_i$. We say that the edge-colored poset P satisfies the *structure condition* for \mathfrak{g} if

$$(9) \quad \text{wt}(x) + \alpha_i = \text{wt}(y) \quad \text{whenever } x \xrightarrow{i} y.$$

We now define two conditions on the pairs of edge labels (c_{yx}, d_{xy}) . We call $\pi_{xy} := c_{yx} d_{xy}$ an edge product. We say that the edge-labeled poset P satisfies the *crossing condition* if for any vertex

y and any color i we have

$$(10) \quad \sum_{x: x \xrightarrow{i} y} \pi_{xy} - \sum_{z: y \xrightarrow{i} z} \pi_{yz} = m_i(y).$$

We say that the edge-labeled poset P satisfies the *diamond condition* if for any pair of vertices (x, y) of the same rank and any pair of colors (i, j) , possibly $i = j$, we have

$$(11) \quad \sum_{z: x \xrightarrow{j} z \text{ and } y \xrightarrow{i} z} c_{zx} d_{yz} = \sum_{u: u \xrightarrow{i} x \text{ and } u \xrightarrow{j} y} d_{ux} c_{yu}.$$

Theorem 3.5. [5, Proposition 3.4] *Given an edge-colored and edge-labeled ranked poset P , the actions (7) and (8) define a representation of \mathfrak{g} on $\mathbb{C}P$ if and only if P satisfies the structure, crossing, and diamond conditions, namely (9), (10), and (11).*

If the given conditions hold, we obtain a *weight basis* of the given representation, and the edge-colored poset P together with its edge-labels is called a *representation diagram*.

Based on representation diagrams, in [13] we give a short proof of the *Gelfand-Tsetlin basis* construction for the irreducible representations of \mathfrak{sl}_n ; recall that this basis is the only one with respect to which the representation matrices of the Chevalley generators are given by explicit formulas. The relevant poset is the set of SSYT of a fixed shape with entries in $[n]$, where an edge colored i corresponds to changing a single entry $i + 1$ to i in a SSYT; this is known as the Gelfand-Tsetlin (distributive) lattice. The edge coefficients are certain rational numbers expressed in a nice factored form in terms of the corresponding SSYT. Based on Theorem 3.5, our proof consists of verifying that these coefficients satisfy (10) and (11).

Similar explicit constructions of certain symplectic and orthogonal irreducible representations (indexed by rows or columns) were given in [4, 7], as well as of the adjoint representations of all simple Lie algebras [6]. Several properties of such bases, such as minimality with respect to the edge set, were also exhibited [5]. However, much more work remains to be done, for instance to extend the mentioned results to an arbitrary symplectic or orthogonal irreducible representation.

4. THE STABLE BEHAVIOR OF SOME GRADED MULTIPLICITIES

This section is based on Richard's paper [37], which studies the decomposition into irreducible components of polynomial functions on the adjoint representation of $G = GL_n(\mathbb{C})$, i.e., the representation on its Lie algebra $\mathfrak{g} = \mathfrak{gl}_n(\mathbb{C})$. (In fact, the setup of the paper is more general, also including a reference to the *q-Dyson conjecture*, and $SL_n(\mathbb{C})$ is used instead.)

The adjoint representation $\text{ad} : G \rightarrow GL(\mathfrak{g})$ is the rational representation defined by $\text{ad}(X)A = XAX^{-1}$. This extends to a representation on the symmetric algebra $S(\mathfrak{g})$, viewed as a graded algebra in the usual way. By a theorem of Kostant, $S(\mathfrak{g})$ is a free module over the ring of G -invariant polynomials $S(\mathfrak{g})^G$, with $S(\mathfrak{g}) \cong S(\mathfrak{g})^G \otimes \mathcal{H}_{\mathfrak{g}}$, where $\mathcal{H}_{\mathfrak{g}}$ are the *G-harmonic polynomials*.

Recall (see, e.g., [44]) that the irreducible rational representations of G are labeled by weakly decreasing sequences of integers with length n or, equivalently, by pairs of partitions $\alpha = (\alpha_1 \geq \dots \geq \alpha_k > 0)$, $\beta = (\beta_1 \geq \dots \geq \beta_l > 0)$ with $k + l \leq n$; the corresponding bijection is given by

$$(\alpha, \beta) \mapsto (\alpha_1, \dots, \alpha_k, 0, \dots, 0, -\beta_l, \dots, -\beta_1) =: [\alpha, \beta]_n.$$

We may decompose $\mathcal{H}_{\mathfrak{g}}$ into irreducible components such that each one is homogeneous. For a fixed pair (α, β) , let d_i be the degrees (also called *generalized exponents*) of the components isomorphic to the irreducible representation labeled by $[\alpha, \beta]_n$, and define

$$G_{[\alpha, \beta]_n}(q) := \sum_i q^{d_i}.$$

Following an idea of Ranee Gupta, who showed that

$$G_{\alpha\beta}(q) := \lim_{n \rightarrow \infty} G_{[\alpha,\beta]_n}(q)$$

exists as a formal power series, Richard expressed the stable graded multiplicities based on the internal product $*$ of Schur functions. This corresponds to the tensor product of irreducible representations of the symmetric group (via the *Frobenius characteristic*). Here we define the internal product formally, in the basis of power sum symmetric functions $p_\lambda := p_{\lambda_1} p_{\lambda_2} \dots = p_1^{m_1} p_2^{m_2} \dots$, by

$$p_\lambda * p_\mu = \delta_{\lambda\mu} z_\lambda p_\lambda, \quad \text{where } z_\lambda := (m_1! 1^{m_1})(m_2! 2^{m_2}) \dots$$

Theorem 4.1. [37][Proposition 8.1] *We have*

$$G_{\alpha\beta}(q) = s_\alpha * s_\beta(q, q^2, \dots).$$

Stembridge [44] developed the combinatorics of the rational representations of GL_n , and used it to express the decomposition of the k th tensor power of \mathfrak{gl}_n (i.e., the adjoint representation) into modules which are simultaneously irreducible with respect to GL_n and the symmetric group S_k . Based on this, he was able to rederive some of Richard's results in [37].

Howe, Tan, and Willenbring [14] showed that Theorem 4.1 is a special case of a general stability result, which we now briefly describe. Let G be a complex reductive algebraic group and $K \subset G$ the fixed-point set of a regular involution on G . Denote by $\mathfrak{k} \subset \mathfrak{g}$ the corresponding Lie algebras, and by \mathfrak{p} the Cartan complement of \mathfrak{k} in \mathfrak{g} . The adjoint action of K on \mathfrak{p} extends to a K -module structure on the symmetric algebra $S(\mathfrak{p})$. By the Kostant-Rallis theorem (which generalizes Kostant's theorem mentioned above), we have $S(\mathfrak{p}) \cong S(\mathfrak{p})^K \otimes \mathcal{H}_\mathfrak{p}$, where the notation is as above. The main result of [14] describes the graded multiplicity of the K -module $\mathcal{H}_\mathfrak{p}$ in the case where (G, K) is a *classical symmetric pair* and the degree d is restricted to the so-called *stable range*. Theorem 4.1 corresponds to $K = GL_n$ diagonally embedded into $G = GL_n \times GL_n$, and the stable range is $n \geq 2d$. There are nine more classical symmetric pairs, such as $(GL_{n+m}, GL_n \times GL_m)$, (SO_{2n}, GL_n) , and (Sp_{2n}, GL_n) , for which the graded multiplicities in the stable range are expressed in terms of the Littlewood-Richardson coefficients, see Section 1.

5. JACK, MACDONALD, AND HALL-LITTLEWOOD SYMMETRIC POLYNOMIALS

In this section I will refer to Richard's paper [41] on *Jack symmetric functions*, as well as to some of the many related developments; indeed, this paper has an impressive number of citations.

5.1. Richard's work on Jack symmetric functions. The Jack functions $J_\lambda^{(\alpha)}(x)$, where λ is a partition, are a remarkable basis of symmetric functions depending on a parameter α . They can be defined as the eigenfunctions of the *generalized Laplace-Beltrami operator*, and for $\alpha = 1$ they specialize (up to normalization) to the Schur functions [28].

Richard's paper [41] is the first one devoted to extending the properties of Schur functions to Jack functions. Some of its results involve: specializations of Jack functions, formulas for special Jack functions and special coefficients, a Cauchy identity (cf. (2)), skew Jack functions and a related coproduct formula, a *Pieri formula* (for the expansion in the basis of Jack functions of the product $J_\lambda^{(\alpha)}(x) J_{(n)}^{(\alpha)}(x)$ by a "one-rowed" Jack function), and a formula for the skew Jack functions in terms of skew SSYT (which is derived from the Pieri formula and generalizes (1)). Richard also states an important conjecture: after suitable renormalization, the structure constants for Jack functions (or analogues of Littlewood-Richardson coefficients for Schur functions) belong to $\mathbb{N}[\alpha]$.

5.2. Recent related developments. In [17], Knop and Sahi prove that the Jack structure constants are in $\mathbb{Z}[\alpha]$, and give a monomial formula for $J_\lambda^{(\alpha)}(x)$ in terms of more general fillings of the Young diagram λ than SSYT (see Section 5.4). The coefficients in this formula are in $\mathbb{N}[\alpha]$, whereas in Richard's formula they are rational functions in α .

In [2], by studying the Laplace-Beltrami operator, Cai and Jing give another monomial formula (with rational function coefficients) for $J_\lambda^{(\alpha)}(x)$, as well as a (non-positive) formula for the Jack structure constants. In another paper, Cai and Jing [3] prove a special case of Richard's conjecture by deriving the generalization of the Pieri rule in which the one-rowed partition is replaced by a rectangle (or a rectangle and another row). Based on this formula, they settle the long-standing problem of realizing any $J_\lambda^{(\alpha)}(x)$ using *vertex operators*.

In [11], Graham and Hunziker give an interesting application of Richard's conjecture to relating tensor products $V_\lambda \otimes V_\mu$ of irreducible representations of a complex reductive algebraic group K to the product of isotypical components $S_\lambda \cdot S_\mu$ in the representation of K on the symmetric algebra of some representation V . In the context of *Hermitian symmetric pairs*, they show that, if V_ν appears in $V_\lambda \otimes V_\mu$, then $S_\nu \subseteq S_\lambda \cdot S_\mu$ in all the classical cases, provided that Richard's conjecture is true.

5.3. Macdonald polynomials. Jack polynomials are specializations of *Macdonald polynomials* $P_\lambda(x_1, \dots, x_n; q, t)$, which are symmetric polynomials depending on two parameters q, t [28]; namely, the former are retrieved by setting $q = t^\alpha$ and letting $t \rightarrow 1$. In fact, the mentioned Macdonald polynomials correspond to the root system of type A_{n-1} , whereas there are versions of them for an arbitrary finite root system. Macdonald polynomials are also eigenfunctions of certain operators.

A breakthrough came with two monomial formulas for Macdonald polynomials: the *Haglund-Haiman-Loehr (HHL) formula* [12] in type A , and the *Ram-Yip formula* [31] in arbitrary type. The former is in terms of certain fillings of the Young diagram λ and is shorter, while the latter is in terms of extensions of the *alcove model* [26, 27], or *LS-gallery model* [10], in the representation theory of Lie algebras. The setup of the Ram-Yip formula allows one to also derive a very general, albeit involved, Littlewood-Richardson multiplication formula [46]. Moreover, it was shown in [46] that, in type A , the Pieri special case of this formula “compresses” to Macdonald's more efficient Pieri formula [28]. This raises the possibility of deriving, in a similar way, other efficient multiplication formulas for Macdonald polynomials and their specializations (such as Jack polynomials).

5.4. Hall-Littlewood polynomials. We briefly explain the compression phenomenon mentioned above and the related setup by referring to the $q = 0$ specialization of Macdonald polynomials, namely to the *Hall-Littlewood polynomials*. The reason we chose this specialization is twofold: more is known about compression, and the description is easier. Nevertheless, we can draw a parallel to Jack polynomials, while a compression result for Macdonald polynomials is given in [23].

We start by explaining the corresponding Ram-Yip formula, which was, in fact, given previously in [30, 33]. As we only refer to type A_{n-1} , we will not use a general root system terminology.

Fix a partition $\lambda = (\lambda_1 \geq \dots \geq \lambda_n = 0)$. The *Bruhat order* \leq on permutations in S_n is defined by its covers: $u < w = u(i, j)$ if $\ell(w) = \ell(u) + 1$, where $\ell(\cdot)$ is the length and (i, j) is a transposition. Let $S_n^\lambda := \{w \in S_n : w(i) < w(i+1) \text{ for } \lambda_i = \lambda_{i+1}\}$. Define the sequence of transpositions

$$\Gamma_k := \Gamma_{k,k} \Gamma_{k,k-1} \dots \Gamma_{k,1}, \quad \text{where } \Gamma_{k,i} = ((i, k+1), (i, k+2), \dots, (i, n)).$$

Define the sequence $\Gamma_\lambda := \Gamma_{\lambda'_1} \Gamma_{\lambda'_2} \dots$, called a λ -*chain*, where λ' is the conjugate of λ . The objects of the alcove model are subwords $T = T_1 T_2 \dots$ of Γ_λ , where T_i is a subword of $\Gamma_{\lambda'_i}$. More precisely, we consider the collection of *admissible pairs* (w, T) , with $w \in S_n$ and $T = (t_1, \dots, t_m)$ as before:

$$(12) \quad \mathcal{A}(\lambda) := \{(w, T) : wt_1 \dots t_{i-1} < wt_1 \dots t_i \text{ for all } i, \text{ and } wT \in S_n^\lambda\}, \text{ where } wT := wt_1 \dots t_m.$$

Define the permutations $\pi_j = \pi_j(w, T) := wT_1 \dots T_j$, where the notation means right multiplication by the transpositions in T_1, \dots, T_j , cf. (12). With (w, T) in $\mathcal{A}(\lambda)$ we associate the filling $\sigma(w, T)$ of λ whose j th column is filled with $\pi_j(1), \dots, \pi_j(\lambda'_i)$ from top to bottom. Clearly, the rows of $\sigma(w, T)$ are weakly increasing, but the columns are not necessarily increasing.

Example 5.1. Let $n = 4$ and $\lambda = (2, 1, 0, 0)$, so $\lambda' = (2, 1)$. We have

$$(13) \quad \Gamma_\lambda = ((2, 3), (2, 4), \underline{(1, 3)}, (1, 4) | (1, 2), (1, 3), \underline{(1, 4)}),$$

where $|$ indicates the concatenation of various $\Gamma_{\lambda'_i}$. Let $T = T_1 T_2$ with $T_1 = ((1, 3))$ and $T_2 = ((1, 4))$, cf. the underlines in (13). Considering w to be the identity in S_4 , we note that $(w, T) \in \mathcal{A}(\lambda)$; indeed, the conditions in (12) are verified (we use the one-line notation for permutations):

$$w = 1234 < w(1, 3) = 3214 = \pi_1 < w(1, 3)(1, 4) = wT = 4213 = \pi_2 \in S_4^\lambda.$$

Hence, the associated filling $\sigma(w, T)$ and the corresponding monomial $x^{\sigma(w, T)}$, cf. (1), are

$$\sigma(w, T) = \begin{array}{|c|c|} \hline 3 & 4 \\ \hline 2 & \\ \hline \end{array}, \quad x^{\sigma(w, T)} = x_2 x_3 x_4.$$

Theorem 5.2. [30, 33] *We have*

$$(14) \quad P_\lambda(x_1, \dots, x_n; t) = \sum_{(w, T) \in \mathcal{A}(\lambda)} t^{\frac{1}{2}(\ell(w) + \ell(wT) - |T|)} (1-t)^{|T|} x^{\sigma(w, T)}.$$

Remark 5.3. By setting $t = 0$ in (14), the only terms that survive correspond to (w, T) for which w is the identity and the chain in Bruhat order in (12) is saturated (i.e., all the steps are covers). The restriction of the map $(w, T) \mapsto \sigma(w, T)$ to these admissible pairs is a bijection to $SSYT_{[n]}(\lambda)$. As it is well-known that $P_\lambda(x_1, \dots, x_n; t = 0)$ is the Schur polynomial $s_\lambda(x_1, \dots, x_n)$, we recover (1). Hence the alcove model gives a new way to realize SSYT as saturated chains in Bruhat order.

We will now state a formula which is essentially the specialization of the HHL formula to $q = 0$. In this sense, it is the analogue of the Knop-Sahi formula for Jack polynomials mentioned above, which is also obtained by specializing the HHL formula, see [12][Section 8].

Let $\mathcal{T}_{[n]}(\lambda)$ denote the fillings σ of λ with entries in $[n]$ satisfying the following conditions: (i) the rows of σ are weakly increasing; (ii) $\sigma(u) \neq \sigma(v)$ whenever the boxes u and v are in the same column or in two consecutive columns, with the right box strictly below the left one (such positions are called *attacking*); (iii) the segment of the leftmost column corresponding to rows $i, i+1, \dots, j$ with $\lambda_i = \lambda_{i+1} = \dots = \lambda_j$ is (strictly) increasing.

Theorem 5.4. [12, 25] *We have*

$$(15) \quad P_\lambda(x_1, \dots, x_n; t) = \sum_{\sigma \in \mathcal{T}_{[n]}(\lambda)} t^{\text{cinv}(\sigma)} (1-t)^{\text{des}(\sigma)} x^\sigma,$$

where $\text{cinv}(\cdot)$ and $\text{des}(\cdot)$ are certain statistics on fillings (see [12, 25]).

There is also an older formula for (skew) Hall-Littlewood polynomials, due to Macdonald [28], which is in terms of (skew) SSYT. This is the analogue of Richard's formula for (skew) Jack polynomials mentioned above. Recall that a tableau $\sigma \in SSYT_{[n]}(\lambda)$ can be represented by a chain of partitions $\lambda^{(0)} \subseteq \lambda^{(1)} \subseteq \dots \subseteq \lambda^{(n)} = \lambda$, where $\lambda^{(i)}$ is the shape of the subtableau consisting of entries $\leq i$. Let $m_i(\lambda) := \lambda'_i - \lambda'_{i+1}$, where λ' is the conjugate partition, as usual.

Theorem 5.5. [28] *We have*

$$(16) \quad P_\lambda(x_1, \dots, x_n; t) = \sum_{\sigma \in SSYT_{[n]}(\lambda)} \frac{\varphi_\sigma(t)}{b_\lambda(t)} x^\sigma,$$

where

$$b_\lambda(t) := \prod_{i \geq 1} \varphi_{m_i(\lambda)}(t), \quad \text{with } \varphi_k(t) := (1-t) \dots (1-t^k), \quad \text{and}$$

$$\varphi_\sigma(t) := \prod_{i=1}^n \varphi_{\lambda^{(i)}/\lambda^{(i-1)}}(t), \quad \text{with } \varphi_{\lambda/\mu}(t) := \prod_{j \in I} 1 - t^{m_j(\mu)} \quad \text{and } I := \{j : \lambda'_j - \mu'_j > \lambda'_{j+1} - \mu'_{j+1}\}.$$

We state a first compression result for Hall-Littlewood polynomials. For simplicity, from now on we assume that λ has no repeated parts, so $S_n^\lambda = S_n$; for the general case, see [25][Section 6].

Theorem 5.6. [25] *The map $(w, T) \mapsto \sigma(w, T)$ is a surjection from $\mathcal{A}(\lambda)$ to $\mathcal{T}_{[n]}(\lambda)$. For all σ in $\mathcal{T}_{[n]}(\lambda)$, the sum in (14) restricted to (w, T) with $\sigma(w, T) = \sigma$ equals the corresponding term in (15).*

We now present a stronger compression for Hall-Littlewood polynomials. In fact, this factors through the first compression, via the surjections $\mathcal{A}(\lambda) \rightarrow \mathcal{T}_{[n]}(\lambda) \rightarrow SSYT_{[n]}(\lambda)$ given by $(w, T) \mapsto \sigma(w, T) \mapsto \bar{\sigma}(w, T)$, where $\bar{\sigma}(w, T)$ is obtained from $\sigma(w, T)$ by sorting columns, cf. Example 5.1.

Theorem 5.7. [16] *For all σ in $SSYT_{[n]}(\lambda)$, the sum in (14) restricted to (w, T) with $\bar{\sigma}(w, T) = \sigma$ equals the corresponding term in (16).*

As stated in Section 5.3, it would be very interesting to derive compressed formulas similar to those in this section for the multiplication structure constants corresponding to Macdonald polynomials or certain specializations of them (e.g., Hall-Littlewood or Jack polynomials). The starting point would be the general Littlewood-Richardson rule in [46] in terms of the alcove model. Generalizing some of these results to other Lie types would also be very useful, cf. [16, 24].

REFERENCES

- [1] G. Benkart and T. Roby. Down-up algebras. *J. Algebra*, 209:305–344, 1998.
- [2] W. Cai and N. Jing. Applications of a Laplace-Beltrami operator for Jack polynomials. *European J. Combin.*, 33:556–571, 2012.
- [3] W. Cai and N. Jing. Jack vertex operators and realization of Jack functions. *J. Algebraic Combin.*, 39:53–74, 2014.
- [4] R. Donnelly. Explicit constructions of the fundamental representations of the symplectic Lie algebras. *J. Algebra*, 233:37–64, 2000.
- [5] R. Donnelly. Extremal properties of bases for representations of semisimple Lie algebras. *J. Algebraic Combin.*, 17:255–282, 2003.
- [6] R. Donnelly. Extremal bases for the adjoint representations of the simple Lie algebras. *Comm. Algebra*, 34:3705–3742, 2006.
- [7] R. Donnelly, S. Lewis, and R. Pervine. Constructions of representations of $\mathfrak{o}(2n + 1, \mathbb{C})$ that imply Molev and Reiner-Stanton lattices are strongly Sperner. *Discrete Math.*, 263:61–79, 2003.
- [8] S. Fomin. Schensted algorithms for dual graded graphs. *J. Algebraic Combin.*, 4:5–45, 1995.
- [9] W. Fulton. *Young Tableaux*, volume 35 of *London Math. Soc. Student Texts*. Cambridge Univ. Press, Cambridge and New York, 1997.
- [10] S. Gaussent and P. Littelmann. LS-galleries, the path model and MV-cycles. *Duke Math. J.*, 127:35–88, 2005.
- [11] W. Graham and M. Hunziker. Multiplication of polynomials on Hermitian symmetric spaces and Littlewood-Richardson coefficients. *Canad. J. Math.*, 61:351–372, 2009.
- [12] J. Haglund, M. Haiman, and N. Loehr. A combinatorial formula for Macdonald polynomials. *J. Amer. Math. Soc.*, 18:735–761, 2005.
- [13] P. Hersh and C. Lenart. Combinatorial constructions of weight bases. The Gelfand-Tsetlin basis. *Electron. J. Combin.*, 17:#R33, 2010.
- [14] R. Howe, E.-C. Tan, and J. Willenbring. The stability of graded multiplicity in the setting of the Kostant-Rallis theorem. *Transform. Groups*, 13:617–636, 2008.
- [15] M. Kashiwara. On crystal bases of the q -analogue of universal enveloping algebras. *Duke Math. J.*, 63:465–516, 1991.
- [16] I. Klostermann. Generalization of the Macdonald formula for Hall-Littlewood polynomials. *Adv. Math.*, 248:1366–1403, 2013.
- [17] F. Knop and S. Sahi. A recursion and a combinatorial formula for Jack polynomials. *Invent. Math.*, 128:9–22, 1997.
- [18] J.-H. Kwon. Demazure crystals of generalized Verma modules and a flagged RSK correspondence. *J. Algebra*, 322:2150–2179, 2009.
- [19] J.-H. Kwon. A plactic algebra of extremal weight crystals and the Cauchy identity for Schur operators. *J. Algebraic Combin.*, 34:427–449, 2011.
- [20] J.-H. Kwon. Crystal bases of q -deformed Kac modules over the quantum superalgebras $U_q(\mathfrak{gl}(m|n))$. *Int. Math. Res. Not.*, (2):512–550, 2014.

- [21] A. Lascoux. Double crystal graphs. In *Studies in memory of Issai Schur (Chevaleret/Rehovot, 2000)*, volume 210 of *Progr. Math.*, pages 95–114. Birkhäuser Boston, Boston, MA, 2003.
- [22] C. Lenart. On the combinatorics of crystal graphs. II. The crystal commutator. *Proc. Amer. Math. Soc.*, 136:825–837, 2008.
- [23] C. Lenart. On combinatorial formulas for Macdonald polynomials. *Adv. Math.*, 220:324–340, 2009.
- [24] C. Lenart. Haglund-Haiman-Loehr type formulas for Hall-Littlewood polynomials of type B and C . *Algebra Number Theory*, 4:887–917, 2010.
- [25] C. Lenart. Hall-Littlewood polynomials, alcove walks, and fillings of Young diagrams (Appendix with A. Lubovsky). *Discrete Math.*, 311:258–275, 2011.
- [26] C. Lenart and A. Postnikov. Affine Weyl groups in K -theory and representation theory. *Int. Math. Res. Not.*, pages 1–65, 2007. Art. ID rnm038.
- [27] C. Lenart and A. Postnikov. A combinatorial model for crystals of Kac-Moody algebras. *Trans. Amer. Math. Soc.*, 360:4349–4381, 2008.
- [28] I. G. Macdonald. *Symmetric Functions and Hall Polynomials*. Oxford Mathematical Monographs. Oxford University Press, Oxford, second edition, 1995.
- [29] R. Proctor. Representations of $\mathfrak{sl}(2, \mathbb{C})$ on posets and the Sperner property. *SIAM J. Algebraic Discrete Methods*, 3:275–280, 1982.
- [30] A. Ram. Alcove walks, Hecke algebras, spherical functions, crystals and column strict tableaux. *Pure Appl. Math. Q.*, 2:963–1013, 2006.
- [31] A. Ram and M. Yip. A combinatorial formula for Macdonald polynomials. *Adv. Math.*, 226:309–331, 2011.
- [32] B. Sagan and R. Stanley. Robinson-Schensted algorithms for skew tableaux. *J. Combin. Theory Ser. A*, 55:161–193, 1990.
- [33] C. Schwer. Galleries, Hall-Littlewood polynomials, and structure constants of the spherical Hecke algebra. *Int. Math. Res. Not.*, pages Art. ID 75395, 31, 2006.
- [34] R. Stanley. Unimodal sequences arising from Lie algebras. In *Combinatorics, representation theory and statistical methods in groups*, volume 57 of *Lecture Notes in Pure and Appl. Math.*, pages 127–136. Dekker, New York, 1980.
- [35] R. Stanley. Weyl groups, the hard Lefschetz theorem, and the Sperner property. *SIAM J. Algebraic Discrete Methods*, 1:168–184, 1980.
- [36] R. Stanley. $GL(n, \mathbb{C})$ for combinatorialists. In *Surveys in combinatorics (Southampton, 1983)*, volume 82 of *London Math. Soc. Lecture Note Ser.*, pages 187–199. Cambridge Univ. Press, Cambridge, 1983.
- [37] R. Stanley. The stable behavior of some characters of $SL(n, \mathbb{C})$. *Linear and Multilinear Algebra*, 16:3–27, 1984.
- [38] R. Stanley. Unimodality and Lie superalgebras. *Stud. Appl. Math.*, 72:263–281, 1985.
- [39] R. Stanley. Differential posets. *J. Amer. Math. Soc.*, 1:919–961, 1988.
- [40] R. Stanley. Log-concave and unimodal sequences in algebra, combinatorics, and geometry. In *Graph theory and its applications: East and West (Jinan, 1986)*, volume 576 of *Ann. New York Acad. Sci.*, pages 500–535. New York Acad. Sci., New York, 1989.
- [41] R. Stanley. Some combinatorial properties of Jack symmetric functions. *Adv. Math.*, 77:76–115, 1989.
- [42] R. Stanley. Variations on differential posets. In *Invariant theory and tableaux (Minneapolis, MN, 1988)*, volume 19 of *IMA Vol. Math. Appl.*, pages 145–165. Springer, New York, 1990.
- [43] R. Stanley. *Enumerative Combinatorics. Vol. 2*, volume 62 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999.
- [44] J. Stembridge. Rational tableaux and the tensor algebra of \mathfrak{gl}_n . *J. Combin. Theory Ser. A*, 46:79–120, 1987.
- [45] M. A. A. van Leeuwen. An analogue of jeu de taquin for Littelmann’s crystal paths. *Sém. Lothar. Combin.*, 41:Art. B41b, 23 pp. (electronic), 1998.
- [46] M. Yip. A Littlewood-Richardson rule for Macdonald polynomials. *Math. Z.*, 272:1259–1290, 2012.

DEPARTMENT OF MATHEMATICS AND STATISTICS, STATE UNIVERSITY OF NEW YORK AT ALBANY, ALBANY, NY 12222, USA

E-mail address: clenart@albany.edu