THE COMBINATORICS OF STEENROD OPERATIONS ON THE COHOMOLOGY OF GRASSMANNIANS

CRISTIAN LENART

Abstract. The study of the action of the Steenrod algebra on the mod p cohomology of spaces has many applications to the topological structure of those spaces. In this paper we present combinatorial formulas for the action of Steenrod operations on the cohomology of Grassmannians, both in the Borel and the Schubert picture. We consider integral lifts of Steenrod operations, which lie in a certain Hopf algebra of differential operators. The latter has been considered recently as a realization of the Landweber-Novikov algebra in complex cobordism theory; it also has connections with the action of the Virasoro algebra on the boson Fock space. Our formulas for Steenrod operations are based on methods which have not been used before in this area, namely Hammond operators and the combinatorics of Schur functions. We also discuss applications of our formulas to the geometry of Grassmannians.

1. Introduction

The action of the *Steenrod algebra* on the mod p cohomology of a space has a deep geometrical significance, whence it offers more information about the geometry of that space than just the cup product structure of its cohomology. Indeed, there are many instances where viewing the cohomology as a module over the Steenrod algebra distinguishes between spaces with isomorphic cohomology rings. Furthermore, the obstruction for maps induced in cohomology to be maps of algebras over the Steenrod algebra is a reasonably strong obstruction for the existence of certain topological maps. The Steenrod algebra can also be used to investigate the attaching maps of CW-complexes, as we discuss in §3 and §6. All these facts show the importance of determining the action of Steenrod operations on the cohomology of spaces we want to study. Among such spaces, the *Grassmannians* play an important role. Apart from projective spaces, very little is known about the attaching maps of their cells.

From a combinatorial point of view, Grassmannians are important because their cohomology ring can be realized naturally in terms of symmetric functions, the Schur functions corresponding to the *Schubert classes*. No good explanation has been found yet for the occurence of Schur functions in both the cohomology of Grassmannians and the representation theory of the symmetric and general linear groups. However, there has been considerable interest recently in connections between the Steenrod algebra and the modular representation theory of the symmetric and general linear groups (see [28], [15], [13], [6]). From this perspective, it is possible that the Steenrod algebra might bring more evidence that the occurence of Schur functions in the two areas mentioned above is not a simple accident, and that there are deep connections between these areas.

In this paper we present formulas for the action of Steenrod operations on the cohomology of Grassmannians, both in the Borel and the Schubert picture. These formulas have applications to the study of the attaching maps of the Schubert cells in the Grassmannians; they might also be relevant in the representation theoretical context discussed above. Our work is based on combinatorial methods which have not been used before in this area, namely Hammond operators and the combinatorics of Schur functions.

In §2, we present a Hopf algebra of differential operators, which has been considered recently by Reg Wood [33] and Nigel Ray [26] as a realization of the Landweber-Novikov algebra in complex cobordism, and which provides integral lifts of Steenrod operations. We point out that the Hopf algebra of symmetric functions has a natural *module Hopf algebra* structure over the algebra of differential operators. Some connections with the action of the *Virasoro algebra* on the *boson Fock* space are also mentioned. In §3, we review the topological background concerning the Steenrod algebra and the cohomology of Grassmannians. In §4, we present an efficient algorithm for computing the action of Steenrod operations on *Chern classes*. This leads to nice generalizations for primes $p > 2$ of the Wu formula (for $p = 2$); previous attempts to generalize it produced only partial results or very complicated closed formulas. In §5, we investigate the action of the integral lifts of Steenrod operations introduced in §2 on Schur functions; this corresponds via reduction mod p to the action of Steenrod operations on Schubert classes in the cohomology of Grassmannians. We give several formulas involving the combinatorics of Young diagrams, analyze their consequences, and conjecture a generalization of one of them. In §6 we apply our formulas in §5 to some simple examples involving Schubert classes in Grassmannians, and suggest possible implications for the corresponding attaching maps.

I am grateful to Nigel Ray and Reg Wood for introducing me to this area, and for pointing out to me the rich combinatorics involved. I am also grateful to Sergey Fomin, Mike Hopkins, Haynes Miller, and Gian-Carlo Rota for valuable discussions.

2. Construction of the Algebra of Differential Operators

In this section we present a purely algebraic construction of certain objects in algebraic topology, and discuss their properties from the same perspective. Any reference to topology is postponed to §3. We refer the reader to [23] for all information concerning Hopf algebras.

Let us first recall the construction of the Hopf algebra $\mathcal{D}^*$ in [33]. R. Wood defines $\mathcal{D}^*$ to be an algebra of differential operators on the polynomial algebra $W^* := \mathbb{Z}[x_1, x_2, \ldots]$, and then uses computations with differential operators to prove a formula for the composition of two basis elements in this algebra. Finally, he dualizes his product formula and obtains the well-known formula for the comultiplication in the dual of the Landweber-Novikov algebra. Here we adopt a different approach, which has the advantage of avoiding the computations with differential operators in [33]. Our starting point is the dual $\mathcal{D}_*$ of the algebra of differential operators, whose comultiplication has a combinatorial description. We then define $\mathcal{D}^*$ by duality, together with a certain action of it on W^*, which has its origins in algebraic topology. We show that this action can be expressed in terms of differential operators, whence the realization of $\mathcal{D}^*$ considered as a definition in [33].

All rings and algebras we consider in this paper are assumed graded by complex dimension, so that products commute without signs.

Let $\mathcal{D}_*$ be the graded polynomial algebra $\mathbb{Z}[b_1, b_2, \ldots]$, where the element b_i has degree i for every i. We consider the formal sum $b := 1+b_1+b_2+\ldots$, and define a comultiplication by

$$\Delta(b_k) = \sum_{i=0}^{k} (b)_{k-i}^{i+1} \otimes b_i \,, \tag{2.1}$$

where $b_0 := 1$ and $(b)_{k-i}^{i+1}$ denotes the part of degree $k-i$ in b^{i+1}. In fact, this amounts to substitution of power series; indeed, if we let $b(t) := t + b_1 t^2 + b_2 t^3 + \ldots$, and we let $b^{(1)}(t)$, $b^{(2)}(t)$ be the corresponding generating functions for $b_k \otimes 1$ and $1 \otimes b_k$, we have

$$\sum_{k\geq 0} \Delta(b_k) t^{k+1} = b^{(2)}(b^{(1)}(t)) \quad \text{in } (\mathcal{D}_* \otimes \mathcal{D}_*)[[t]] \,. \tag{2.2}$$

It is not difficult to show that $\mathcal{D}_*$ is a non-cocommutative Hopf algebra. Indeed, it is the *incidence Hopf algebra* corresponding to the family of compositions of integers, which are Boolean algebras (see [27]). The counit is specified by

$$\varepsilon(b_k) = \begin{cases} 1 & \text{if } k = 0 \\ 0 & \text{otherwise} \,, \end{cases}$$

for all $k \geq 0$, and the antipode is given by Lagrange inversion; more precisely, $S(b_k)$ is the coefficient of t^{k+1} in the substitutional inverse of $b(t)$, that is the coefficient of t^k in

$$\frac{1}{k+1} \left(\frac{t}{b(t)} \right)^{k+1} .$$

Let us consider the graded dual $\mathcal{D}^*$ of $\mathcal{D}_*$. Given a partition $\lambda \vdash n$, we denote by D_λ the element of $\mathcal{D}^*$ which is dual to $b_\lambda := b_{\lambda_1} b_{\lambda_2} \ldots$ with respect to the monomial basis of $\mathcal{D}_*$. We abbreviate $D_{(k)}$ to D_k. In this paper, we use the standard notation for partitions of positive integers, namely given $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_l) = (1^{m_1} 2^{m_2} \ldots k^{m_k})$ with $\lambda_1 \geq \lambda_2 \geq \ldots \geq \lambda_l > 0$, we write λ' for its *conjugate*, and

$$l(\lambda) := l \,, \quad |\lambda| := \lambda_1 + \ldots + \lambda_l \,, \quad \|\lambda\| := m_1! \ldots m_k! \,;$$

recall that $l(\lambda)$ is known as the length of λ, and $|\lambda|$ as the weight of λ. Clearly, the degree of D_λ is $|\lambda|$. The comultiplication in $\mathcal{D}^*$ can be easily obtained by duality, and is given by

$$\Delta(D_\lambda) = \sum_{\mu \cup \nu = \lambda} D_\mu \otimes D_\nu \,,$$

where the union of partitions is defined by concatenating them and reordering the parts. The multiplication in $\mathcal{D}^*$ will be denoted by $\circ$. R. Wood's formula for composing two basis elements D_λ and D_μ (see [33]) can be obtained directly from (2.1) by duality.

We now define a left action of $\mathcal{D}^*$ on the graded polynomial algebra $W^* := \mathbb{Z}[x_1, x_2, \ldots]$ (every x_i has degree 1). First of all, we let D_0 (the unit in $\mathcal{D}^*$) act as the identity, and we define

$$D_\lambda \, 1 = 0 \,, \quad D_\lambda \, x_i := \begin{cases} x_i^{k+1} & \text{if } \lambda = (k) \\ 0 & \text{otherwise} \,, \end{cases} \tag{2.3}$$

for every $\lambda \vdash k \neq 0$. We then define the action of D_λ on a product by the so-called Cartan formula

$$D_\lambda (yz) = \sum_{\mu \cup \nu = \lambda} (D_\mu\, y)\, (D_\nu\, z)\,. \tag{2.4}$$

The fact that this is a good definition follows from the coassociativity and the cocommutativity of the comultiplication in $\mathcal{D}^*$. However, it is not obvious that the formulas above define a left action. The following theorem contains the proof of this fact.

Proposition 2.5. *W^* is a left $\mathcal{D}^*$-*module algebra*, i.e. the multiplication and the unit maps of W^* are $\mathcal{D}^*$-module maps.*

Proof. All we have to prove is that formulas (2.3) and (2.4) define a left action of $\mathcal{D}^*$ on W^*. Indeed, once we know this, the two formulas tell us precisely that W^* is a $\mathcal{D}^*$-module algebra.

We need to show that

$$(D_\lambda \circ D_\mu)\, y = D_\lambda\, (D_\mu\, y)\,, \tag{2.6}$$

for every $y \in W^*$, and every partitions $\lambda \vdash m$, $\mu \vdash k$. Let us first consider the case $y = x_i$. Assume for the moment that $\mu = (k)$ and $l(\lambda) \leq k+1$. By (2.3) and (2.4), the RHS of (2.6) is αx_i^{m+k+1}, where α is the number of terms of the form $D_{i_1} \otimes \ldots \otimes D_{i_{k+1}}$ in the k-fold comultiplication $\Delta_k(D_\lambda)$. But this is just the number of permutations of the multiset consisting of the parts of λ and $k+1-l(\lambda)$ zeros, which is

$$\frac{(k+1)!}{\|\lambda\|\,(k+1-l(\lambda))!}\,.$$

On the other hand, $(D_\lambda \circ D_\mu)\, x_i$ is βx_i^{m+k+1}, where

$$\beta = \langle D_\lambda \circ D_k \mid b_{m+k} \rangle = \langle D_\lambda \otimes D_k \mid \Delta(b_{m+k}) \rangle\,.$$

By (2.2), β is the coefficient of $b_\lambda t^{m+k+1}$ in $b(t)^{k+1}$, so the multinomial theorem gives $\beta = \alpha$. If $l(\mu) > 1$ or $l(\lambda) > k+1$, the above argument shows that both sides of (2.6) are 0. Finally, if y is a certain monomial in the x_i's, a straightforward induction on its degree based on the Cartan formula (2.4) shows that (2.6) holds in general. □

We now intend to realize the algebra $\mathcal{D}^*$ as an algebra of differential operators. Let us recall some notation from [33]. The *Weyl algebra* $\mathcal{W}^*$ is the associative algebra with unit (over the rationals) which is generated by x_i, ∂_j (i, j non-negative integers) subject to the relations

$$[x_i, x_j] = 0\,,\ \ [\partial_i, \partial_j] = 0\,,\ \ [x_i, \partial_j] = \delta_{i,j}\,;$$

here the square brackets denote the Lie product, and δ_{ij} the Kronecker delta. We let $x_\lambda := x_{\lambda_1} x_{\lambda_2} \ldots$, and similarly for ∂_λ. There is a natural action of the Weyl algebra on W^* (∂_i acts as partial derivative with respect to x_i), and a natural grading ($x_\lambda\, \partial_\mu$ has degree $l(\lambda) - l(\mu)$). The Weyl algebras $\mathcal{W}_n$ are defined for each n in a similar way, by restricting to the finite set of variables $x_1, \ldots, x_n$ and the corresponding partial derivatives. For every integer k, we consider the summands $\mathcal{W}_n^k$ of degree k, the inverse limit $\widehat{\mathcal{W}}^k$ with respect to the restriction maps, and $\widehat{\mathcal{W}}^* := \bigoplus_k \widehat{\mathcal{W}}^k$. Clearly, the action of $\mathcal{W}^*$ on W^* extends to an action of $\widehat{\mathcal{W}}^*$. We let $\widetilde{D}_0$ be the unit of $\widehat{\mathcal{W}}^*$, and consider the element

$D_k := \sum_{i\geq 1} x_i^{k+1}\partial_i$ in $\widehat{\mathcal{W}}^k$ for every $k \geq 1$. A new multiplication, denoted by $\vee$, is defined on $\widehat{\mathcal{W}}^*$ in [33] by

$$x_\lambda \partial_\mu \vee x_\nu \partial_\pi := x_\lambda x_\nu \partial_\mu \partial_\pi \,;$$

this is a good definition, because every element of $\widehat{\mathcal{W}}^*$ can be written uniquely in a form (known as the *standard form*) where the polynomial part is on the left, and the derivatives are on the right. For every partition $\lambda = (1^{m_1} 2^{m_2} \dots k^{m_k})$, we define

$$D_\lambda := \frac{D_1^{\vee m_1}}{m_1!} \vee \ldots \vee \frac{D_k^{\vee m_k}}{m_k!}\,,$$

where $D_i^{\vee n}$ denotes the n-fold $\vee$-multiplication. We intend to show that we can identify the operator D_λ in $\mathcal{D}^*$ with the operator D_λ, whence we can identify $\mathcal{D}^* \otimes \mathbb{Q}$ with the corresponding subalgebra of $\widehat{\mathcal{W}}^*$.

Theorem 2.7. *The $\mathbb{Z}$-module spanned by D_λ is closed under composition in $\widehat{\mathcal{W}}^*$, and can be identified with $\mathcal{D}^*$ as a graded algebra.*

Proof. It is not difficult to show that every linear map from W^* to W^{*+k} corresponds to the action of a unique element of $\widehat{\mathcal{W}}^*$, which obviously has degree k. Hence all we need to prove is that the actions of D_λ and D_λ on W^* coincide. Let $\lambda = (1^{m_1} 2^{m_2} \dots k^{m_k})$. Clearly, the actions of the two operators on every x_i coincide. Hence it suffices to prove that D_λ also satisfies the Cartan formula (2.4).

We consider matrices A with k rows and an infinite number of columns, which have non-negative integer entries a_{ij}. We write $r(A) = (r_1(A), \dots, r_k(A))$ and $c(A) = (c_1(A), c_2(A), \dots)$ for the row and column sums of A, assuming that the former are finite. Given such a matrix A and a sequence $a = (a_1, a_2, \dots)$, we also use the following notation

$$x_A := \prod_{j\geq 1} x_j^{2a_{1j}+\ldots+(k+1)a_{kj}}\,, \quad \partial_a := \prod_{j\geq 1} \partial_j^{a_j}\,, \quad \binom{r(A)}{A} := \prod_{i=1}^{k} \binom{r_i(A)}{a_{i1}, a_{i2}, \dots};$$

note that the multinomial coefficients are well-defined, since the matrix A has only a finite number of non-zero elements in each row. Given the sequence $m = (m_1, \dots, m_k)$ considered above, we use the multinomial theorem to compute

$$\|\lambda\| D_\lambda = \sum_{r(M)=m} \binom{m}{M} x_M\, \partial_{c(M)}\,. \tag{2.8}$$

Combining this formula with the Leibniz formula, we have that

$$(\|\lambda\| D_\lambda)(yz) = \sum_{r(M)=m} \sum_{s+t=c(M)} \left(\prod_{j\geq 1} \binom{c_j(M)}{s_j}\right) \binom{m}{M} x_M\, (\partial_s\, y)\, (\partial_t\, z)\,.$$

On the other hand, using (2.8) once again, we have

$$\|\lambda\| \sum_{\mu\cup\nu=\lambda} (D_\mu\, y)(D_\nu\, z) = \sum_{p+q=m} \sum_{\substack{r(P)=p\\ r(Q)=q}} \left(\prod_{i=1}^{k} \binom{m_i}{p_i}\right) \binom{p}{P} \binom{q}{Q} x_{P+Q}\, (\partial_{c(P)}\, y)\, (\partial_{c(Q)}\, z)\,.$$

In order to show that the left-hand sides of the last two formulas coincide, it is enough to show that the corresponding coefficients of $x_M\, (\partial_s\, y)\, (\partial_t\, z)$ are equal for every matrix M

with $r(M) = m$, and every infinite sequences of positive integers s, t with $s + t = c(M)$. This amounts to proving the following identity:

$$\sum_{\substack{P+Q=M \\ c(P)=s,\, c(Q)=t}} \left(\prod_{i=1}^{k} \binom{m_i}{r_i(P)} \right) \binom{r(P)}{P} \binom{r(Q)}{Q} = \left(\prod_{j \geq 1} \binom{c_j(M)}{s_j} \right) \binom{m}{M} .$$

This clearly holds, since both sides count distributions of integers into boxes B_{ij}, $i = 1, \ldots, k$, $j \geq 1$ such that: (1) the i-th row of boxes contains the numbers $1, \ldots, m_i$ with a plus or minus sign attached to them; (2) box B_{ij} contains m_{ij} elements; (3) there are s_j positive numbers in the j-th column of boxes. □

From now on we drop the notation D_λ and think of D_λ as the corresponding differential operator (as it is in fact defined [33]).

Remarks 2.9.

(1) The $\vee$-multiplication in $\mathcal{D}^*$ is the transpose of the comultiplication in $\mathcal{D}_*$ specified by $b_k \mapsto 1 \otimes b_k + b_k \otimes 1$. This shows that $\mathcal{D}^*$ is also a Hopf algebra with respect to the $\vee$-multiplication and the same comultiplication. In fact, it is a tensor product of divided power Hopf algebras in D_k, for $k \geq 1$.
(2) It is known from [33] that the algebra $\mathcal{D}^* \otimes \mathbb{Q}$ is generated under composition by D_1 and D_2 only.
(3) It is easy to check that the Lie bracket of D_k and D_l is given by $[D_k, D_l] = (l - k)D_{k+l}$. Therefore, the $\mathbb{Z}$-module spanned by D_k is a Lie algebra, and its rational universal enveloping algebra is $\mathcal{D}^* \otimes \mathbb{Q}$.
(4) D_λ maps W^n (that is polynomials of degree n) to $W^{n+|\lambda|}$. Hence the action map $\mathcal{D}^* \otimes W^* \to W^*$ is a map of graded algebras.
(5) We have that $D_\lambda \, x_\mu = 0$ whenever $l(\lambda) > l(\mu)$.
(6) The action of $\mathcal{D}^*$ on W^* is clearly faithful. On the other hand, the space of invariants of this action consists of 0 only; this can be easily seen by considering the action of D_k on a homogeneous polynomial y, and observing that the leading monomial of $D_k \, y$ can be expressed in terms of the leading monomial of y.

Let us now consider the algebra of symmetric functions with integer coefficients in $x_1, x_2, \ldots$, which we denote by Sym_*. We use the notation of [19] for symmetric functions, namely h_λ for the complete homogeneous symmetric functions, e_λ for the elementary symmetric functions, p_λ for the power sums, m_λ for the monomial symmetric functions, and s_λ for the Schur functions. It is well-known that Sym_* is a polynomial algebra (over the integers) in both h_n and e_n, and that $Sym_* \otimes \mathbb{Q}$ is a polynomial algebra (over the rationals) in p_n. In fact, Sym_* is a Hopf algebra in which p_n are primitives, and

$$\delta(h_n) = \sum_{k=0}^{n} h_k \otimes h_{n-k} \, , \quad \delta(e_n) = \sum_{k=0}^{n} e_k \otimes e_{n-k} \, . \tag{2.10}$$

The graded dual Sym^* is isomorphic to Sym_*, and we have the standard pairing $\langle m_\lambda | h_\mu \rangle = \delta_{\lambda\mu}$. The transpose of the multiplication by a symmetric function f is denoted by $f^\perp$, and the corresponding operator is known as a *Hammond operator*. For instance $p_n^\perp = n \, \partial / \partial p_n$. As pointed out in [19], the linear span of p_n and $p_n^\perp$ is a Heisenberg Lie algebra.

The action of $\mathcal{D}^*$ on W^* can be extended to an action on the power series ring $\widehat{W}_* := \mathbb{Z}[[x_1, x_2, \ldots]]$, and then restricted to a left action on Sym^*. By partial duality, we obtain a right action of $\mathcal{D}^*$ on Sym_*, which we denote by $D^\perp f$ for every f in Sym_* and D in $\mathcal{D}^*$. On the other hand, if we consider duality rather than partial duality, we obtain a left coaction of $\mathcal{D}_*$ on Sym_*. In fact, there are two more actions and three more coactions that can be obtained in a similar way, and any of them determines all the others (see [2]).

According to Proposition 2.5, Sym^* is a left $\mathcal{D}^*$-module algebra. We now show that in fact it is a *module Hopf algebra*, which means that the comultiplication and the antipode are also $\mathcal{D}^*$-module maps.

Theorem 2.11. *Sym^* is a left $\mathcal{D}^*$-module Hopf algebra, and Sym_* is a right $\mathcal{D}^*$-module Hopf algebra. On the other hand, Sym_* is a left $\mathcal{D}_*$-comodule Hopf algebra.*

Proof. We need to check that the so-called co-Cartan formula holds:

$$\delta(D\, f) = (\Delta(D))\,(\delta(f))\,, \tag{2.12}$$

where D is an arbitrary element in $\mathcal{D}^*$, $f \in Sym^*$, Δ is the comultiplication in $\mathcal{D}^*$, and δ is the comultiplication in Sym^*. This can be easily checked for $f = p_n$ and $D = D_\lambda$ using the fact that $D_\lambda\, p_n = \alpha\, p_{n+|\lambda|}$ for a certain integer α (cf. the computation in the proof of Proposition 2.5). On the other hand, it is also easy to show that if (2.12) holds for the symmetric functions f_1 and f_2, it holds for $f_1 f_2$, by using the Cartan formula (2.4). A similar argument applies to the antipode. The other statements can be obtained by partial duality and duality, respectively. □

Remarks 2.13.

(1) The condition for the comultiplication in Sym^* to be a $\mathcal{D}^*$-module map is equivalent to the map $\mathcal{D}^* \otimes Sym^* \to Sym^*$ being a coalgebra map.
(2) Similarly, the condition for the multiplication in Sym_* to be a $\mathcal{D}_*$-comodule map is equivalent to the coaction map $Sym_* \to \mathcal{D}_* \otimes Sym_*$ being an algebra map.

The coaction of $\mathcal{D}_*$ on Sym_* is also related to substitution of power series. In order to express it, we introduce the generating function for the complete homogeneous symmetric functions $h(t) := 1 + h_1 t + h_2 t^2 + \ldots$; we also let $h^{(2)}(t)$ be the corresponding generating function for $1 \otimes h_k$ (note the difference from the definition of $b(t)$).

Proposition 2.14. *The coaction map $Sym_* \to \mathcal{D}_* \otimes Sym_*$ (which we observed that is an algebra map) is specified by*

$$h(t) \mapsto h^{(2)}(b^{(1)}(t))\,.$$

Proof. Let us denote by $\widetilde{\Delta}$ the coaction map. The coefficient of $b_\lambda \otimes h_\mu$ in $\widetilde{\Delta}(h_n)$ is

$$\alpha_{\lambda\mu} = \langle D_\lambda \otimes m_\mu \mid \widetilde{\Delta}(h_n)\rangle = \langle D_\lambda\, m_\mu \mid h_n\rangle\,,$$

where $|\lambda| + |\mu| = n$. By (2.8), this coefficient is 0 unless $l(\mu) = 1$. So let $\mu = (k)$, where $k := n - |\lambda|$. By the computation in the proof of Proposition 2.5, we have

$$\alpha_{\lambda(k)} = \begin{cases} \frac{k!}{\|\lambda\|(k-l(\lambda))!} & \text{if } l(\lambda) \le k \\ 0 & \text{otherwise}\,. \end{cases}$$

But by the multinomial theorem, this is exactly the coefficient of $b_\lambda\, t^{|\lambda|+k}$ in $b(t)^k$, which concludes the proof. □

Finally, we show that the actions of the operators D_k and $D_k^\perp$ on Sym^* and Sym_* are closely related to the action of the *Virasoro algebra* on the *boson Fock space*.

Proposition 2.15. *For every $k \geq 1$, the operators D_k and $D_k^\perp$ on Sym^* and Sym_* can be expressed as*

$$D_k = \sum_{i \geq 1} p_{i+k}\, p_i^\perp\,, \quad D_k^\perp = \sum_{i \geq 1} p_i\, p_{i+k}^\perp\,.$$

Proof. The actions of D_k and $\sum_{i\geq 1} p_{i+k}\, p_i^\perp$ clearly agree on p_j, since $D_k\, p_j = j p_{j+k}$. On the other hand, both operators are derivations, whence their actions agree on any symmetric function. The proof of the statement concerning $D_k^\perp$ is similar. □

We briefly recall some basic concepts from conformal field theory in order to discuss the connection with the algebra $\mathcal{D}^*$; for more details, the reader is referred to [12], for instance. The boson Fock space of central charge 1 is defined to be

$$\mathcal{H}_T := \mathbb{Q}[p_1, p_2, \ldots][u, u^{-1}]\,,$$

where p_k can be thought of as the power sum symmetric functions. The charge n sector of $\mathcal{H}_T$ is $\mathbb{Q}[p_1, p_2, \ldots]u^n$. The Virasoro algebra of central charge c is the Lie algebra over $\mathbb{Q}$ spanned by elements L_k, $k \in \mathbb{Z}$, which has the following bracket:

$$[L_k, L_l] = (k-l)L_{k+l} + \frac{k^3 - k}{12}\, c\, \delta_{k,-l}\,.$$

There is a natural action of the Virasoro algebra on the boson Fock space. For $c = 1$, the basis elements of the Virasoro algebra act on the charge 0 sector of $\mathcal{H}_T$ in the following way:

$$L_0 = \sum_{i \geq 1} p_i\, p_i^\perp\,,$$

$$L_k = \sum_{i \geq 1} p_i\, p_{i+k}^\perp + \frac{1}{2} \sum_{i=1}^{k-1} p_i^\perp\, p_{k-i}^\perp\,,$$

$$L_{-k} = \sum_{i \geq 1} p_{i+k}\, p_i^\perp + \frac{1}{2} \sum_{i=1}^{k-1} p_i\, p_{k-i}\,,$$

where $k \geq 1$. We immediately observe that the expressions for D_k and $D_k^\perp$ in Proposition 2.15 are just truncated versions of the expressions for L_{-k} and L_k, respectively (in fact $D_1 = L_{-1}$ and $D_1^\perp = L_1$). Furthermore, the Lie algebras spanned by $\{D_k : k \geq 0\}$ and $\{D_k^\perp : k \geq 0\}$ are isomorphic to the corresponding Lie subalgebras of the Virasoro algebra. However, the elements D_k and $D_k^\perp$ considered together do not span a Lie algebra.

We conclude by mentioning that T. Katsura et al. [14] constructed a natural embedding of the dual of the Landweber-Novikov algebra (which is just the Hopf algebra $\mathcal{D}_*$, as shown in the next section) in the charge 0 sector of the boson Fock space.

3. Background from Algebraic Topology

In this section, we briefly review the topology underlying the algebra presented in §2. We refer the reader to [30] and [21] for all information concerning the Steenrod algebra, to [1] for information about the Landweber-Novikov algebra, and to [10] or [11] for information about the cohomology of Grassmannians.

For every prime p, the mod p *Steenrod algebra* $\mathcal{A}(p)$ is the graded associative algebra of mod p stable operations in ordinary cohomology theory over $\mathbb{F}_p$. If $p = 2$, it is generated by the Steenrod squares Sq^n ($n \geq 1$), modulo the Adem relations; if p is odd, it is generated by the Bockstein operation and the (reduced) Steenrod operations $\mathcal{P}^n$ ($n \geq 1$), modulo the corresponding Adem relations. The operations Sq^n raise degree by n, while the operations $\mathcal{P}^n$ raise degree by $2n(p-1)$; in other words:

$$Sq^n \colon H^k(-, \mathbb{F}_p) \to H^{k+n}(-, \mathbb{F}_p)\,, \quad \mathcal{P}^n \colon H^k(-, \mathbb{F}_p) \to H^{k+2n(p-1)}(-, \mathbb{F}_p)\,.$$

For simplicity, we will denote Sq^{2n} by $\mathcal{P}^n$ as well. Milnor showed that $\mathcal{A}(p)$ modulo the two-sided ideal generated by the Bockstein operation has a linear basis indexed by compositions of positive integers. Furthermore, he showed that both $\mathcal{A}(p)$ and its dual have a natural Hopf algebra structure over $\mathbb{F}_p$. The action of the Steenrod algebra on the cohomology of spaces gives rise (by partial duality) to an action of it on homology, and to a coaction of the dual Steenrod algebra on homology (by duality). There are two more actions and three more coactions that can be obtained in a similar way, and they are discussed in [2].

There are many generalized cohomology theories now used in algebraic topology, and most of them have a Hopf algebra of stable operations analogous to the Steenrod algebra. In every such situation, there is a geometric way to define four actions and four coactions of these algebras or their duals on the corresponding cohomology or homology. An important such cohomology theory is complex cobordism $MU^*(-)$ introduced by Milnor in [22]. The structure of the algebra $MU^*(MU)$ of all operations in complex cobordism was determined by Landweber [18] and Novikov [24]. They showed that $MU^*(MU)$ is isomorphic to the tensor product of the complex cobordism ring MU_* and the so-called *Landweber-Novikov algebra*, which turns out to be precisely the Hopf algebra $\mathcal{D}^*$ discussed in the previous section; in fact $MU^*(MU)$ is a so-called *Hopf algebroid*.

Let us now refer to the Grassmannian of n-planes in $\mathbb{C}^\infty$, which is also the classifying space for principle $U(n)$-bundles, whence the notation $BU(n)$; note that $BU(1)$ is the infinite complex projective space, usually denoted by $\mathbb{C}P^\infty$. There is a natural inclusion of $BU(n)$ into $BU(n+1)$, so we can consider the union of all $BU(n)$, which is the space denoted by BU. Let $E^*(-)$ be an unreduced multiplicative cohomology theory with complex orientation $Z \in E^2(\mathbb{C}P^\infty)$; in particular, $E^*(-)$ can be ordinary cohomology (with integer or $\mathbb{F}_p$ coefficients) or complex cobordism. We write $E_* := E_*(\text{point})$ for the coefficient ring, and identify E_n with $E^{-n} := E^{-n}(\text{point})$, as it is usually done. It is well-known that $E^*(BU(n)) \cong E^*[c_1, c_2, \ldots, c_n]$ and $E^*(BU) \cong E^*[c_1, c_2, \ldots]$, where c_i are the generalized *Chern classes*. This description of the cohomology of $BU(n)$ is called the *Borel picture*. It is also known that the map

$$E^*(BU(n)) \to E^*(\mathbb{C}P^\infty \times \ldots \times \mathbb{C}P^\infty) \cong E^*[x_1, x_2, \ldots, x_n]$$

induced by the classifying map of the direct product of n copies of the Hopf bundle over $\mathbb{C}P^\infty$ is a monomorphism mapping c_i, with $i \leq n$, to the i-th elementary symmetric polynomial in $x_1 := Z \otimes 1 \otimes \ldots \otimes 1$, $x_2 := 1 \otimes Z \otimes \ldots \otimes 1$, ..., $x_n := 1 \otimes 1 \otimes \ldots \otimes Z$. On the other hand, we have that $E_*(BU) \cong E_*[b_1, b_2, \ldots]$, and that c_n is dual to b_1^n with respect to the monomial basis of $E_*[b_1, b_2, \ldots]$. The multiplicative structure of $E_*(BU)$ is determined by the map $BU \times BU \to BU$ classifying the Whitney sum of vector bundles. The diagonal map $BU \to BU \times BU$ induces a comultiplication $\delta \colon E_*(BU) \to E_*(BU \times BU) \cong E_*(BU) \otimes E_*(BU)$ satisfying

$$\delta(b_n) = \sum_{i=0}^{n} b_i \otimes b_{n-i} \, ,$$

which turns $E_*(BU)$ into a Hopf algebra. It follows from the above considerations that we may identify $E_*(BU)$ with $Sym_* \otimes E_*$ and $E^*(BU)$ with $Sym^* \otimes E^*$, in such a way that b_n is identified with h_n and c_n with e_n.

We can also consider the Grassmannian $Gr_n(\mathbb{C}^{n+k})$ of n-planes in $\mathbb{C}^{n+k}$, whose cohomology ring is a certain quotient of the cohomology of $BU(n)$; indeed, the obvious inclusion $Gr_n(\mathbb{C}^{n+k}) \hookrightarrow BU(n)$ induces the quotient map. Let λ be an arbitrary partition (identified with its Young diagram) with at most n rows and k columns, and let $F_\bullet$ be a fixed complete flag

$$0 = F_0 \subset F_1 \subset F_2 \subset \ldots \subset F_{n+k} = \mathbb{C}^{n+k}$$

of subspaces with $\dim(F_i) = i$. There is a *Schubert cell* $\Omega_\lambda = \Omega_\lambda(F_\bullet)$ defined by

$$\Omega_\lambda := \{V \in Gr_n(\mathbb{C}^{n+k}) \, : \, \dim(V \cap F_{i+\lambda_{n+1-i}}) \geq i \, , \;\; 1 \leq i \leq n\} \, ;$$

here we set $\lambda_i = 0$ for $i > l(\lambda)$. It is known that Ω_λ is an irreducible closed subvariety of $Gr_n(\mathbb{C}^{n+k})$ of dimension $2|\lambda|$. It is also known that $\Omega_\mu \subset \Omega_\lambda$ if and only if the Young diagram of μ is strictly contained in that of λ, and that Ω_λ is the disjoint union of the interiors of the cells Ω_μ, $\mu \subseteq \lambda$. Furthermore, the cohomology class σ_λ of Ω_λ (that is, the dual of the fundamental class of Ω_λ) is independent of the choice of the fixed flag defining it. These classes, known as Schubert classes, form a basis over the integers for the cohomology ring of $Gr_n(\mathbb{C}^{n+k})$. This description of the cohomology of $Gr_n(\mathbb{C}^{n+k})$ is called the *Schubert picture*. The crucial fact is that the quotient map from the cohomology of $BU(n)$ (which we identified with the ring of symmetric polynomials in n variables) to the cohomology ring of $Gr_n(\mathbb{C}^{n+k})$ maps the Schur polynomial $s_\lambda(x_1, \ldots, x_n)$ to σ_λ if the Young diagram λ has at most n rows and k columns, and to 0 otherwise.

Another topological fact we need concerns the relation between the action of Steenrod operations on Schubert cells and the corresponding attaching maps. It follows from the naturality of Steenrod operations that any composition of them applied to a Schubert class is a linear combination (with non-zero coefficients) of Schubert classes whose corresponding cells are non-trivially attached to the initial cell. Hence, whenever we detect a non-zero coefficient, we can say that we have a non-trivial attaching map. Conversely, when every composition of Steenrod operations applied to a Schubert class gives a zero coefficient for a higher dimensional Schubert class, we have a chance that the corresponding attaching map is trivial.

Let us now consider the infinite product of infinite complex projective spaces, whose E^*-cohomology is $W^* \otimes E^* = E^*[x_1, x_2, \ldots]$. The action of the Landweber-Novikov algebra on $MU^*[x_1, x_2, \ldots]$ restricts to an action on W^*, which coincides with the action of the algebra $\mathcal{D}^*$ discussed in §2. This was proved by N. Ray [26] by comparing the action of the operators D_λ with the action of the Landweber-Novikov operations described, for instance, in [1]. On the other hand, the action of the Steenrod operations on $\mathbb{F}_p[x_1, x_2, \ldots]$ is given by the following rules:

(1) $\mathcal{P}^0$ is the identity;
(2) $\mathcal{P}^1 x_i = x_i^p$ and $\mathcal{P}^n x_i = 0$ for $n > 1$ $(i \geq 1)$;
(3) $\mathcal{P}^n (yz) = \sum_{k=0}^{n} (\mathcal{P}^k y)(\mathcal{P}^{n-k} z)$ for every y, z in $W^* \otimes \mathbb{F}_p$ (Cartan formula).

In fact, the Cartan formula holds for every space, while $\mathcal{P}^n$ raises every cohomology class x to its p-th power if $\dim(x) = 2n$, and maps it to 0 if $\dim(x) < 2n$. This justifies the name "reduced power operations" (or "squaring operations" for $p = 2$). As far as the action on the cohomology of BU is concerned, we have the following formula due to Borel and Serre [3]:

$$\mathcal{P}^n c_{k+n} = m_{(1^k p^n)}\,; \tag{3.1}$$

here m_λ is the corresponding monomial symmetric function in $H^*(BU, \mathbb{F}_p)$, identified with the ring of symmetric functions with $\mathbb{F}_p$ coefficients. This formula was used in [17] to lift the action of $\mathcal{P}^n$ on the mod p cohomology of BU to the integral cohomology. It is not difficult to see that the operator defining this integral lift is precisely the differential operator $D_{((p-1)^n)}$ in $\mathcal{D}^*$. In fact, as the three rules above show, this operator lifts the action of $\mathcal{P}^n$ on the mod p cohomology of the infinite product $\mathbb{C}P^\infty \times \mathbb{C}P^\infty \times \ldots$ to the integral cohomology. Furthermore, the subalgebra of $\mathcal{D}^*$ generated by the operators $D_{((p-1)^n)}$ $(n \geq 0)$ under composition can be identified upon reduction mod p with the quotient of $\mathcal{A}(p)$ by the two-sided ideal generated by the Bockstein operation. This is proved in [33] for $p = 2$, but the argument can be easily extended to any prime. Let us mention R. Wood's notation SQ^n for $D_{(1^n)}$, which we will also use.

4. Steenrod Operations in the Borel Picture of the Cohomology of Grassmannians

In this section we present an efficient algorithm for computing the action of Steenrod operations on Chern classes in $H^*(BU, \mathbb{F}_p)$. This problem has been studied for a long time. According to (3.1), it is just the problem of expanding the monomial symmetric function $m_{(1^k p^n)}$ in the basis of elementary symmetric functions. We can ask for this expansion over the integers, which would correspond to the integral lift of Steenrod operations, or we could just ask for the mod p reduction. In principle, we can then compute the action of Steenrod operations on any monomial in the Chern classes using the Cartan formula.

For $p = 2$, the mod 2 formula was found by Wu [34]:

$$m_{(1^k 2^n)} = \sum_{i=0}^{n} \binom{n+k-i-1}{n-i} e_i\, e_{2n+k-i} \pmod 2\,. \tag{4.1}$$

The corresponding integral formula was given by Carlitz [7]. As far as the case $p > 2$ is concerned, it is shown in [17] that the coefficient of e_{pn+k} in $m_{(1^k p^n)}$ is $\frac{pn+k}{n+k}\binom{n+k}{n}$. This agrees with the mod p result of Brown and Peterson [5], who show that the coefficient is $\binom{n+k-1}{n}$ mod p. Other partial results appeared in [4] and [25]. On the other hand, Shay [29] obtains a closed formula for the expansion of $m_{(1^k p^n)}$ in the elementary symmetric functions by using the expansion in the power sums as an intermediate step. However, his formula is complicated and difficult to work with; indeed, it is a non-trivial matter to deduce from this formula when a given coefficient is 0.

The general problem of expanding a monomial symmetric function m_λ in the basis of elementary ones goes back to Cayley and MacMahon [20]. Their idea is to apply the Hammond operators $h_i^\perp$ to the expansion of m_λ with indeterminate coefficients. It is easy to see that $h_k^\perp$ acts on a monomial symmetric function by removing a part equal to k, if it exists, and gives 0 otherwise; furthermore, $h_k^\perp$ satisfies the Cartan formula. Hence we obtain a system of equations which depends on the expansion of monomial symmetric functions indexed by partitions obtained from λ by removing one part. In this section, we discuss a refinement of this algorithm which is specially designed for the case $\lambda = (1^k p^n)$, and is based on symmetric functions in two sets of variables. Let us also note that there are combinatorial formulas for the expansion of an arbitrary monomial symmetric function in terms of the lattice of partitions of a set [9], as well as in terms of certain combinatorial objects called *layered primitive bi-brick permutations* [16]. However, both of these formulas are complicated and difficult to work with, even in the special case we consider.

For the rest of this section, we fix an arbitrary integer $p \geq 2$ (not necessarily a prime). We use the following notation:

$$m_{(1^k p^n)} = \sum_{\lambda \vdash pn+k} \alpha_\lambda^{n,k} \, e_\lambda \, . \tag{4.2}$$

We also denote by $\lambda - 1$ the partition obtained from λ by subtracting 1 from every part. The following Proposition collects a couple of simple observations.

Proposition 4.3.

(1) *We have* $\alpha_\lambda^{n,k} = 0$ *unless* $\lambda \geq (n^{p-1}(n+k))$ *in the dominance order; furthermore,* $\alpha_{(n^{p-1}(n+k))}^{n,k} = 1$. *In particular,* $\alpha_\lambda^{n,k} = 0$ *unless* $l(\lambda) \leq p$ *and* $\lambda_1 \geq n+k$.
(2) *If* $l(\lambda) = p$, *then* $\alpha_\lambda^{n,k} = \alpha_{\lambda-1}^{n-1,k}$.

Proof. 1) This result is in fact more general. We show first that for any λ, we have

$$e_\lambda = m_{\lambda'} + \sum_{\mu < \lambda'} \gamma_{\lambda\mu} \, m_\mu \, .$$

It is easy to see that the partitions μ occuring in the above sum are obtained in the following way: we place λ'_1 boxes in a column, then add λ'_2 boxes such that no two are in the same row, then continue in the same way. Clearly, $\mu \leq \lambda'$ in the dominance order. It is also easy to see that the coefficient of $m_{\lambda'}$ is 1. We now use the above relation and induction on λ to show that

$$m_\lambda = e_{\lambda'} + \sum_{\nu > \lambda'} \theta_{\lambda\nu} \, e_\nu \, .$$

Here we also need the well-known fact that $\mu < \lambda$ if and only if $\mu' > \lambda'$.

2) This follows easily by applying the Hammond operator $h_p^\perp$ to both sides of (4.2). □

According to the previous observations, it suffices to determine the coefficients $\alpha_\lambda^{n,k}$ corresponding to partitions λ of length at most $p-1$. The crucial ingredient for our algorithm is working with symmetric functions in two sets of variables $x = (x_1, x_2, \ldots)$ and $y = (y_1, y_2, \ldots)$. Let

$$h_n := h_n(x)\,, \quad \overline{h}_n := h_n(y)\,, \quad H_n := h_n(x,y)\,,$$

and similarly for the elementary symmetric functions. There are corresponding Hammond operators $H_n^\perp$, and they also satisfy the Cartan formula

$$H_n^\perp\,(uv) = \sum_{k=0}^{n} (H_k^\perp\, u)\,(H_{n-k}^\perp\, v)\,.$$

It is known that $H_n = \sum_{k=0}^n h_k\,\overline{h}_{n-k}$. Let us also note that $H_1^\perp\, e_n = e_{n-1}$ and $H_1^\perp\,\overline{e}_n = \overline{e}_{n-1}$.

Now let us fix n. We define $\overline{m}_{(1^k p^n)}$ in $\mathbb{Z}[x,y]$ as follows:

$$\overline{m}_{(1^k p^n)} := \begin{cases} \sum_{\lambda \vdash pn+k} \alpha_\lambda^{n,k}\, e_{\lambda\setminus\lambda_1}\,\overline{e}_{\lambda_1} & \text{if } k \geq n(p-2) \\ \\ (H_1^\perp)^{n(p-2)-k}\,\overline{m}_{(1^{n(p-2)}p^n)} & \text{otherwise}\,; \end{cases}$$

the notation $\lambda \setminus i$ means the partition obtained from λ by removing the part i. Clearly, $\overline{m}_{(1^k p^n)}$ has the following form:

$$\overline{m}_{(1^k p^n)} = \sum_\lambda \beta_\lambda^{n,k}\, e_\lambda\,\overline{e}_{pn+k-|\lambda|}\,.$$

Lemma 4.4. *We have $H_1^\perp\,\overline{m}_{(1^k p^n)} = \overline{m}_{(1^{k-1}p^n)}$ for $k \geq 1$, $H_p^\perp\,\overline{m}_{(1^k p^n)} = \overline{m}_{(1^k p^{n-1})}$ for $n \geq 1$, and $H_i^\perp\,\overline{m}_{(1^k p^n)} = 0$ for $i \notin \{1, p\}$.*

Proof. To prove the first formula, it is enough to consider the case $k > n(p-2)$. By using Proposition 4.3, we deduce that in this case $\lambda_1 > \lambda_2$ for every partition λ with at least two parts for which $\alpha_\lambda^{n,k} \neq 0$. Indeed, on the one hand we have $n + k > np - n$, and on the other hand we have $\lambda_1 \geq n + k$, which implies $\lambda_2 \leq np - n$. We conclude that the terms of $H_1^\perp\,\overline{m}_{(1^k p^n)}$ have the form $\beta\, e_{\lambda\setminus\lambda_1}\,\overline{e}_{\lambda_1}$. The first formula now follows by combining this remark with the definition of $\overline{m}_{(1^{k-1}p^n)}$ and the fact that $H_1^\perp$ lifts the action of $h_1^\perp$ on Sym_* to the ring $\mathbb{Z}[e_1, e_2, \ldots; \overline{e}_1, \overline{e}_2, \ldots]$. The other two formulas follow in a similar way if $k > n(p-2)$. Otherwise, we use the commutativity of the operators $H_i^\perp$:

$$\begin{aligned} H_p^\perp\,\overline{m}_{(1^k p^n)} &= H_p^\perp((H_1^\perp)^{n(p-2)-k+1}\,\overline{m}_{(1^{n(p-2)+1}p^n)}) = (H_1^\perp)^{n(p-2)-k+1}(H_p^\perp\,\overline{m}_{(1^{n(p-2)+1}p^n)}) \\ &= (H_1^\perp)^{n(p-2)-k+1}\,\overline{m}_{(1^{n(p-2)+1}p^{n-1})} = \overline{m}_{(1^k p^{n-1})}\,; \end{aligned}$$

a similar argument applies to $H_i^\perp$ with $i \notin \{1, p\}$. □

The following Theorem contains the results underlying our algorithm.

Theorem 4.5.

(1) *The following recurrence relation holds for all* λ *and* $n, k \geq 1$*:*

$$\beta_\lambda^{n,k} - \beta_\lambda^{n,k-1} = (-1)^{p-1} \beta_\lambda^{n-1,k} .$$

(2) *We have* $\beta_\lambda^{n,k} = 0$ *whenever* $l(\lambda) > p-1$ *or* $|\lambda| > n(p-1)$*. Furthermore,* $\beta_\lambda^{n,0} = 0$ *(mod* p*) if* $n > \lambda_1$*.*

(3) *We have* $\alpha_\lambda^{n,k} = \sum_i \beta_{\lambda \setminus i}^{n,k}$*, where the summation ranges over all distinct parts of* λ*. If* $n + k > \lambda_1$*, then* $\alpha_{\lambda \cup (pn+k-|\lambda|)}^{n,k} = \beta_\lambda^{n,k}$*.*

Proof. 1) We apply the operators $H_i^\perp$, $i = 1, 2, \ldots, p$ to $\overline{m}_{(1^k p^n)}$, and identify the coefficient of $e_\lambda \overline{e}_{pn+k-i}$ in the result. By Lemma 4.4, we have

$$\begin{aligned}
\beta_\lambda^{n,k-1} &= \beta_\lambda^{n,k} + \sum_{\lambda^{(1)}} c_{\lambda^{(1)}} \beta_{\lambda^{(1)}}^{n,k} , \\
0 &= \sum_{\lambda^{(1)}} c_{\lambda^{(1)}} \beta_{\lambda^{(1)}}^{n,k} + \sum_{\lambda^{(2)}} c_{\lambda^{(2)}} \beta_{\lambda^{(2)}}^{n,k} , \\
&\ldots \\
0 &= \sum_{\lambda^{(p-2)}} c_{\lambda^{(p-2)}} \beta_{\lambda^{(p-2)}}^{n,k} + \beta_{\lambda^{(p-1)}}^{n,k} , \\
\beta_\lambda^{n-1,k} &= \beta_{\lambda^{(p-1)}}^{n,k} .
\end{aligned}$$

Here $\lambda^{(i)}$ ranges over partitions with at most $p-1$ parts obtained by adding i boxes to the Young diagram of λ, such that no two of them are on the same row; $c_{\lambda^{(i)}}$ are certain positive integer coefficients which we do not specify. The recurrence relation for $\beta_\lambda^{n,k}$ now follows by taking the alternating sum of the formulas above.

2) Clearly, if $\beta_\lambda^{n,k} \neq 0$, then $l(\lambda) \leq p-1$ by Proposition 4.3. If $k \geq n(p-2)$, then $|\lambda| > n(p-1)$ implies $\beta_\lambda^{n,k} = 0$, by the same Proposition; the definition of $\overline{m}_{(1^k p^n)}$ now shows that this is true in general. If $n > \lambda_1$, then $\beta_\lambda^{n,0} = \alpha_{\lambda \cup (pn-|\lambda|)}^{n,0}$ (by assertion 3). But $\alpha_{\lambda \cup (pn-|\lambda|)}^{n,0} = 0 \pmod p$, since $e_n^p = m_{(p^n)} \pmod p$.

3) This formula follows immediately from the fact that $H_1^\perp$ lifts the action of $h_1^\perp$ on Sym_* to the ring $\mathbb{Z}[e_1, e_2, \ldots; \overline{e}_1, \overline{e}_2, \ldots]$. If $n + k > \lambda_1$, then $\beta_{(\lambda \setminus i) \cup (pn+k-|\lambda|)}^{n,k} = 0$ for all parts i of λ (by assertion 2). Hence, in this case $\alpha_{\lambda \cup (pn+k-|\lambda|)}^{n,k} = \beta_\lambda^{n,k}$. □

The reason we are working in $\mathbb{Z}[e_1, e_2, \ldots; \overline{e}_1, \overline{e}_2, \ldots]$ is that the recurrence relation in Theorem 4.5 (1) does not hold for the coefficients $\alpha_\lambda^{n,k}$. The main advantage of our method is that we can express the coefficient $\alpha_\lambda^{n,k} \mod p$ in a very nice form, as the following Corollary shows. Here, as well as throughout this paper, we adopt the usual conventions $\binom{i}{j} = 0$ if $j > i$ or $j < 0$.

Corollary 4.6. *Let* p *be a prime,* λ *a partition with at most* $p-1$ *parts, and* $n_0 := \lceil |\lambda|/(p-1) \rceil$*,* $n_1 := \lambda_1$*. For all* $n, k \geq 0$*, we can express* $\beta_\lambda^{n,k} \mod p$ *in the following way:*

$$\sum_{i=n_0}^{n_1} \beta_\lambda^{i,0} \binom{n+k-i-1}{n-i} .$$

Proof. By the recurrence relation in Theorem 4.5 (1), we have

$$(-1)^{p-1}\beta_\lambda^{n,k} = \beta_\lambda^{n,0} + \sum_{j=1}^{k} \beta_\lambda^{n-1,j}\,.$$

The expression for $\beta_\lambda^{n,k}$ immediately follows by induction using the fact that $\beta_\lambda^{n,0} = 0 \pmod p$ unless $n_0 \le n \le n_1$ (cf. Theorem 4.5 (2)), and the iterated Pascal formula

$$\sum_{j=1}^{k} \binom{q+j-1}{q} = \binom{q+k}{q+1}\,.$$

□

Note the similarity between our expression for $\beta_\lambda^{n,k} \bmod p$ and the Wu formula. For $p = 2$ we have $\beta_i^{n,0} = 0$ if $n < i$, and $\beta_i^{n,0} = 1$ if $n = i$; otherwise, we have $\beta_i^{n,0} = 0 \pmod 2$. The Corollary gives $\beta_i^{n,k} = \binom{n+k-i-1}{n-i}$. But $\alpha_{(2n+k-i,i)}^{n,k} = \beta_i^{n,k}$ if we assume $2n+k-i \ge i$; indeed, if strict inequality holds, we have $2n+k-i > n$, so $\beta_{2n+k-i}^{n,k} = 0$. Hence we obtain the Wu formula (4.1). Thus, our Corollary turns out to be the natural generalization of this formula to primes $p > 2$.

We now give recurrence relations for the coefficients $\beta_\lambda^{n,0}$, so that we have a complete description of an algorithm for expanding $m_{(1^k p^n)}$ in the basis of elementary symmetric functions. If we are interested in the expansion over the integers, we need all $\beta_\lambda^{n,0}$ for $n \ge n_0$, but if we are working mod p (as is the case with Steenrod operations), we only need $\beta_\lambda^{n,0}$ for $n_0 \le n \le n_1$. We then determine $\beta_\lambda^{n,k}$ for $k > 0$, and $\alpha_\lambda^{n,k}$ as a certain sum of the former (cf. Theorem 4.5 (3)). Let us introduce the following notation: given two partitions $\lambda = (1^{l_1} \dots k^{l_k})$ and $\mu = (1^{m_1} \dots k^{m_k})$, we let

$$c_{\lambda\mu} := \binom{m_{k-1}}{l_k - m_k} \binom{m_{k-2}}{l_{k-1} + l_k - m_{k-1} - m_k} \cdots \binom{m_1}{l_2 + \ldots + l_k - m_2 - \ldots - m_k}\,.$$

Proposition 4.7. *For every partition λ, we have $\beta_\lambda^{n,0} = \beta_{\lambda-1}^{n-1,0}$ if $l(\lambda) = p-1$, and*

$$\sum_\mu c_{\lambda\mu}\,\beta_\mu^{n,0} + \sum_\nu c_{\lambda\nu}\,\beta_\nu^{n,0} = 0\,;$$

here μ ranges over partitions obtained from λ by decrementing at most $p-1-l(\lambda)$ parts different from 1 by 1 and adding as many parts equal to 1; on the other hand, ν ranges over partitions obtained from λ by decrementing $k \le p-2-l(\lambda)$ parts different from 1 by 1 and adding $k+1$ parts equal to 1.

Proof. The recurrence relation follows easily by applying the operator $H_{l(\lambda)+1}^{\perp}$ to $\overline{m}_{(1^k p^n)}$ and identifying the coefficient of $e_{\lambda-1}\,\overline{e}_{pn+k-|\lambda|-1}$ in the result. The coefficient $c_{\lambda\mu}$ represents the number of ways to obtain the partition $\lambda - 1$ from μ by subtracting 1 from certain parts of μ, and then reordering the parts (see the diagram below, where $\lambda = (1^{l_1} \dots k^{l_k})$ and $\mu = (1^{m_1} \dots k^{m_k})$; the black boxes are removed from λ, and the shaded ones are added in order to obtain μ).



□

Note that $c_{\lambda\lambda} = 1$, and that the above recurrence relations express $\beta_\lambda^{n,0}$ with $l(\lambda) < p-1$ in terms of $\beta_\mu^{n,0}$ with $l(\mu) > l(\lambda)$.

Example 4.8. We can use the above results to give a closed mod p formula for $\alpha_\lambda^{n,k}$ in the case $p = 3$. A similar formula was obtained in [31], by using the method in [29].

It follows from the discussion above that it suffices to consider partitions λ with two parts. In other words, it suffices to compute $\beta_i^{n,k}$, which in turn depend on $\beta_i^{n,0}$. We know that $\beta_i^{n,0} = 0$ if $i > 2n$, and that $\beta_i^{n,0} = 0 \pmod 3$ if $n > i$. By Proposition 4.7, we have $\beta_i^{n,0} = -\beta_{i-1}^{n-1,0} - \beta_{i-2}^{n-1,0}$ (we let $\beta_j^{n,0} = 0$ if $j \leq 0$). Hence $\beta_i^{n,0}$ mod 3 is completely determined by $\beta_1^{1,0} = -1$ and $\beta_2^{1,0} = -2$ (see the discussion in the next paragraph). A straightforward iteration of the recurrence relation gives

$$\beta_i^{n,0} = (-1)^j \sum_{l=0}^{j} \binom{j}{l} \beta_{i-j-l}^{n-j,0} .$$

Letting $j = n-1$, we obtain $\beta_i^{n,0} = (-1)^{n-1} \frac{2i}{n} \binom{n}{i-n} \pmod 3$. Combining this formula with Corollary 4.6 gives

$$\beta_i^{n,k} = \sum_{j=\lceil i/2 \rceil}^{i} (-1)^{j-1} \frac{2i}{j} \binom{j}{i-j} \binom{n+k-j-1}{n-j} \pmod 3 . \tag{4.9}$$

Finally, assuming that $3n + k \geq 2i$, we have

$$\alpha_{(3n+k-i,i)}^{n,k} = \begin{cases} \beta_i^{n,k} & \text{if } n+k > i \\ \beta_i^{n,k} + \beta_{3n+k-i}^{n,k} & \text{otherwise} . \end{cases} \tag{4.10}$$

The same method can be used to compute $\beta_i^{n,k}$ mod p for any prime $p > 2$. All we have to show is that $\beta_i^{1,0} = -i$ for $1 \leq i \leq p-1$. The easiest way to see this is to note that $\beta_i^{1,0} = \alpha_{(p,i)}^{1,i}$ for $1 \leq i \leq p-1$, which is the same as the coefficient of $e_p e_1^i$ in the expansion of $m_{(p,i)} = m_p m_i - m_{p+i}$ (by the symmetry of the transition matrix from the basis $\{m_\lambda\}$ to $\{e_\lambda\}$); then a simple application of the Waring formula gives the answer. Everything else follows exactly in the same way as above.

Example 4.11. Let us compute the coefficient of $e_1^2 e_2 e_{5n+k-4}$ in $m_{(1^k 5^n)}$ for all n, k with $5n + k > 4$.

For $n = 1$, we have $\beta^{1,0}_{(1^2 2)} + 4\beta^{1,0}_{(1^4)} + \beta^{1,0}_{(1^3 2)} = 0$, whence $\beta^{1,0}_{(1^2 2)} = -4$. On the other hand, $\beta^{1,0}_{(1^3)} + \beta^{1,0}_{(1^4)} = 0$, whence $\beta^{1,0}_{(1^3)} = -1$. In consequence,

$$\alpha^{1,0}_{(1^3 2)} = \beta^{1,0}_{(1^2 2)} + \beta^{1,0}_{(1^3)} = -5\,, \quad \alpha^{1,k}_{(1^2 2(k+1))} = \beta^{1,0}_{(1^2 2)} = -4\,, \;\; k \geq 1\,.$$

For $n = 2$, we have to compute first $\beta^{2,0}_{(1^2 2)} + 4\beta^{2,0}_{(1^4)} + \beta^{2,0}_{(1^3 2)} = 0$. But $\beta^{2,0}_{(1^4)} = \beta^{1,0}_{\emptyset} = 5$ (the coefficient of e_5 in m_5), and $\beta^{2,0}_{(1^3 2)} = \beta^{1,0}_{1} = -1$, by the observation at the end of Example 4.8. Hence $\beta^{2,0}_{(1^2 2)} = -19$. Similarly, we obtain $\beta^{2,0}_{(1^2 6)} = 4$. In consequence,

$$\alpha^{2,0}_{(1^2 26)} = \beta^{2,0}_{(1^2 2)} + \beta^{2,0}_{(1^2 6)} = -15\,, \quad \alpha^{2,k}_{(1^2 2(k+6))} = -19 - 4k\,, \;\; k \geq 1\,.$$

For $n \geq 3$, we only have to compute $\beta^{n,0}_{(1^2 2)} = \alpha^{n,0}_{(1^2 2(5n-4))}$. If we are only interested in working mod 5, then we can apply Corollary 4.6 and obtain

$$\alpha^{n,k}_{(1^2 2(5n+k-4))} = \frac{2n+k-3}{n-1}\binom{n+k-3}{n-2} \pmod 5\,, \qquad \text{for } n \geq 3,\, k \geq 0\,.$$

5. Steenrod Operations in the Schubert Picture of the Cohomology of Grassmannians

In this section, we investigate the action of the operators $D_{(k^n)}$ in $\mathcal{D}^*$ (in particular, SQ^n) on Schur functions. As we have pointed out in §3, this corresponds to the action of Steenrod operations on Schubert classes in the cohomology of Grassmannians. More precisely, for every prime p, the mod p reduction of the action of $D_{((p-1)^n)}$ gives the action of $\mathcal{P}^n$; in particular, the mod 2 reduction of the action of SQ^n gives the action of Sq^{2n}.

Let us start by recalling some terminology concerning Young diagrams. The *content* $c(b)$ of a box b in row $i(b)$ and column $j(b)$ is the difference $j(b) - i(b)$. The *hooklength* $h(b)$ is defined to be $\lambda_{i(b)} + \lambda'_{j(b)} - i(b) - j(b) + 1$. A shape λ is called a *border strip* if it is connected and does not contain a 2×2 block of boxes. A shape satisfying just the second condition is called a *broken border strip*; clearly, such a shape is a union of connected components, each of which is a border strip. The *height* $\mathrm{ht}(\lambda)$ of a broken border strip λ is defined to be one less than the number of rows it occupies. A *sharp corner* in a broken border strip is a box having no boxes above it or to its left. A *dull corner* in a broken border strip is a box that has a box to its left and a box above it, but no box directly northwest of it. We denote by $\mathrm{SC}(\lambda)$ and $\mathrm{DC}(\lambda)$ the sets of sharp and dull corners of the broken border strip λ, and by $cc(\lambda)$ the number of its connected components; if the skew shape λ is not a broken border strip, it is convenient to set $cc(\lambda) := \infty$.

Proposition 5.1. *We have that*

$$D_k\, s_\lambda = \sum_{\substack{\mu \supset \lambda\,:\, |\mu| - |\lambda| = k \\ cc(\mu/\lambda) \leq 2}} d_{\lambda\mu}\, s_\mu\,, \qquad D_k^\perp\, s_\lambda = \sum_{\substack{\mu \subset \lambda\,:\, |\lambda| - |\mu| = k \\ cc(\lambda/\mu) \leq 2}} d_{\mu\lambda}\, s_\mu\,;$$

if μ/λ is a border strip, then

$$d_{\lambda\mu} = (-1)^{\mathrm{ht}(\mu/\lambda)} \left(\sum_{b \in \mathrm{SC}(\mu/\lambda)} c(b) - \sum_{b \in \mathrm{DC}(\mu/\lambda)} c(b) \right),$$

and if μ/λ has two connected components, then $d_{\lambda\mu} = (-1)^{\mathrm{ht}(\mu/\lambda)}$.

Proof. We start by recalling the following formula (see [19]):

$$p_k \, s_\lambda = \sum_{\substack{\mu \supset \lambda \, : \, |\mu| - |\lambda| = k \\ cc(\mu/\lambda) = 1}} (-1)^{\mathrm{ht}(\mu/\lambda)} \, s_\mu \, . \tag{5.2}$$

By duality, we have a similar formula:

$$p_k^{\perp} \, s_\lambda = \sum_{\substack{\mu \subset \lambda \, : \, |\lambda| - |\mu| = k \\ cc(\lambda/\mu) = 1}} (-1)^{\mathrm{ht}(\lambda/\mu)} \, s_\mu \, . \tag{5.3}$$

We now combine these formulas with the expression for D_k in Proposition 2.15. We deduce that $D_k \, s_\lambda$ is a linear combination of s_μ, where three cases are possible: (1) both $\pi := \mu/\lambda$ and $\rho := \lambda/\mu$ are border strips; (2) λ is contained in μ and $\pi := \mu/\lambda$ is a border strip; (3) λ is contained in μ and $\pi := \mu/\lambda$ is a broken border strip with two connected components. These cases are illustrated in the three figures below, where the black boxes are removed, the bold ones are added, and the shaded ones are first removed (by applying the operator $p_i^{\perp}$), and then added (by applying the operator p_{k+i}).



In the first case, let $\theta \subset \lambda$ be the border strip "connecting" π and ρ, i.e., the smallest border strip for which $\pi \cup \theta \cup \rho$ is connected (the shaded area in the first figure). The partition μ can be obtained in two ways from λ: by removing ρ and adding π, or by removing $\rho \cup \theta$ and adding $\pi \cup \theta$. Note that $\mathrm{ht}(\pi \cup \theta) = \mathrm{ht}(\pi) + \mathrm{ht}(\theta) + 1$ and $\mathrm{ht}(\rho \cup \theta) = \mathrm{ht}(\rho) + \mathrm{ht}(\theta)$ if π is southwest of ρ, and viceversa, otherwise. Hence, the corresponding coefficients given by the two formulas above cancel.

In the second case, the partition μ can be obtained in several ways, by removing a border strip θ for which $\pi \cup \theta$ is connected, and then adding $\pi \cup \theta$. The border strip θ can be situated northeast of π, in which case $\mathrm{ht}(\pi \cup \theta) = \mathrm{ht}(\pi) + \mathrm{ht}(\theta) + 1$, or southwest of π, in which case $\mathrm{ht}(\pi \cup \theta) = \mathrm{ht}(\pi) + \mathrm{ht}(\theta)$. In the first case, the sum of the corresponding coefficients is $(-1)^{\mathrm{ht}(\pi)+1}(i(b_1)-1)$, while in the second case, the sum is $(-1)^{\mathrm{ht}(\pi)}(j(b_2)-1)$;

here b_1 and b_2 are the first and last sharp corners of π (from northeast to southwest). On the other hand, it is easy to see that

$$\sum_{b\in \mathrm{SC}(\pi)} c(b) - \sum_{b\in \mathrm{DC}(\pi)} c(b) = \left(\sum_{b\in \mathrm{SC}(\pi)} j(b) - \sum_{b\in \mathrm{DC}(\pi)} j(b) \right) - \left(\sum_{b\in \mathrm{SC}(\pi)} i(b) - \sum_{b\in \mathrm{DC}(\pi)} i(b) \right) = j(b_2) - i(b_1)\,.$$

In the third case, there is a unique way to obtain μ from λ, namely by removing the border strip θ "connecting" π, and then adding $\pi \cup \theta$. Since $\mathrm{ht}(\pi \cup \theta) = \mathrm{ht}(\pi) + \mathrm{ht}(\theta)$, we have $d_{\lambda\mu} = (-1)^{\mathrm{ht}(\pi)}$, as claimed. □

Remark 5.4. Proposition 5.1 has applications to the inverse Kostka matrix $K^{(-1)}$ (see [19]). For example, we can immediately deduce the following closed formula (appearing in [19] p. 107) for $K^{(-1)}_{\lambda\mu}$ with $\lambda = (1^k n)$, $n > 1$:

$$K^{(-1)}_{\lambda\mu} = \begin{cases} (-1)^{n+1}\,(k+1) & \text{if } \mu = (1^{k+n}) \\ (-1)^{n-l} & \text{if } \mu = (1^i 2^j l)\,,\ \ j+l \le n \\ 0 & \text{otherwise}\,; \end{cases}$$

indeed, we only need to notice that $D_{n-1}\, e_{k+1} = m_{(1^k n)}$, which means that in this case $K^{(-1)}_{\lambda\mu} = d_{(1^{k+1})\mu}$ for all μ. In fact, we can use this idea more generally, and give an efficient algorithm for computing the inverse Kostka matrix. For every partition λ, we define an operator M_λ in $\mathcal{D}^*$ recursively as follows:

$$M_k := D_{k-1}\,, \quad M_\lambda := D_{\lambda_1 - 1}\, M_{\lambda\setminus\lambda_1} - \sum_{i=2}^{l(\lambda)} \lambda_i\, M_{(\lambda\setminus\lambda_1\setminus\lambda_i)\cup(\lambda_1+\lambda_i-1)} \ \text{ for } l(\lambda) \ge 2\,.$$

It is easy to see by induction that $M_\lambda\, e_{l(\lambda)} = m_\lambda$. On the other hand, we can express $M_\lambda\, e_{l(\lambda)}$ in the basis of Schur functions using the formula in Proposition 5.1 repeatedly. Thus, we determine $K^{(-1)}_{\lambda\mu}$ for all partitions μ.

In order to express the action of $D_{(k^n)}$, we need the following identity in $\mathcal{D}^*$:

$$n\, D_{(k^n)} = \sum_{i=1}^{n} (-k-1)^{i-1}\, D_{ik} \circ D_{(k^{n-i})}\,. \tag{5.5}$$

R. Wood proves this identity in [33] for $k = 1$, using his composition formula for two basis elements in $\mathcal{D}^*$; his argument can be extended without difficulty to any k. By combining this recursive formula for $D_{(k^n)}$ with Proposition 5.1, we obtain the following Corollary.

Corollary 5.6. *We have that*

$$D_{(k^n)}\, s_\lambda = \sum_{\mu\supset\lambda\,:\,|\mu|-|\lambda|=kn} a_{\lambda\mu}\, s_\mu\,;$$

the coefficients $a_{\lambda\mu}$ are given by

$$a_{\lambda\mu} = \sum_{\lambda=\mu_{(0)}\subset\mu_{(1)}\subset\ldots\subset\mu_{(m)}=\mu} f(\mu_{(0)},\mu_{(1)})\, f(\mu_{(1)},\mu_{(2)})\, \ldots\, f(\mu_{(m-1)},\mu_{(m)})\,;$$

here $\mu_{(i)}/\mu_{(i-1)}$ is a broken border strip with at most two connected components whose length is a multiple of k for every i, and

$$f(\pi,\rho) = \frac{k\,(-k-1)^{(|\rho|-|\pi|)/k-1}}{|\rho|-|\lambda|}\,d_{\pi\rho}\,.$$

Remark 5.7. Since Sq^{2n} squares every cohomology class of dimension $2n$, the coefficient $a_{\lambda\mu}$ for $k=1$ and $|\mu|=2|\lambda|$ is congruent mod 2 to the Littlewood-Richardson coefficient $c^{\mu}_{\lambda\lambda}$. Thus all our results for $a_{\lambda\mu}$ (Proposition 5.6, Theorem 5.12, Corollaries 5.20 and 5.21) can be translated in terms of mod 2 congruences for $c^{\mu}_{\lambda\lambda}$.

Corrolary 5.6 has an important consequence concerning the way in which the coefficients $a_{\lambda\mu}$ depend on the embedding of the skew diagram μ/λ and its connected components in the plane. In order to state this fact, we need additional notation. Let $\pi=\pi(\lambda,\mu)$ and $\rho=\rho(\lambda,\mu)$ be the diagrams of smallest weight (π possibly empty) for which $\mu/\lambda=\rho/\pi$, and let $c(\lambda,\mu)$ be the content of the top left box of $\rho(\lambda,\mu)$ (in its embedding in μ). Similarly, we write the connected components of μ/λ (from northeast to southwest say) as $\rho_{(i)}(\lambda,\mu)/\pi_{(i)}(\lambda,\mu)$, $i=1,2,\ldots$, where the corresponding Young diagrams are again of minimum weight. The corresponding contents are denoted by $c_i(\lambda,\mu)$.

Corollary 5.8. *The coefficient $a_{\lambda\mu}$ depends only on the connected components of μ/λ and the corresponding contents $c_i(\lambda,\mu)$. In particular, it depends only on μ/λ and $c(\lambda,\mu)$.*

Proof. According to Proposition 5.1, the coefficient $d_{\lambda\mu}$ depends on $c(\lambda,\mu)$ if and only if μ/λ is connected. The Corollary follows by induction on $|\mu|-|\lambda|$ by combining the previous observation with (5.5). □

From now on we concentrate on the case $k=1$. In other words we work with the operators $SQ^n = D_{(1^n)}$, which reduce to the Steenrod squares Sq^{2n} upon reduction mod 2. In order to investigate the way in which $a_{\lambda\mu}$ depends on $c(\lambda,\mu)$, we need the following two lemmas.

Lemma 5.9. *The operator $\sum_{i\geq 1} p_i\,p_i^{\perp}$ multiplies every homogeneous symmetric function by its degree. In particular, we have the following generalization of Newton's identity (which can be recovered by choosing $\lambda=(1^n)$):*

$$|\lambda|\,s_\lambda = \sum_{\mu\subset\lambda\,:\,cc(\lambda/\mu)=1} (-1)^{\mathrm{ht}(\lambda/\mu)}\,p_{|\lambda|-|\mu|}\,s_\mu\,.$$

Proof. Using the same technique as in the proof of Proposition 2.15, we can show that the operator $\sum_{i\geq 1} p_i\,p_i^{\perp}$ coincides with the operator $\sum_{i\geq 1} x_i\,\partial_i$. We can easily check that the latter multiplies every monomial in the x_i's by its degree. The identity follows by combining this result with identity (5.3). □

Lemma 5.10. *For every symmetric functions f,g, and every $k>0$, we have*

$$f^{\perp}\,(p_k g) = p_k\,(f^{\perp}\,g) + (p_k^{\perp}\,f)^{\perp}\,g\,.$$

In particular, we have the following generalization of (5.2):

$$p_k\, s_{\mu/\lambda} = \sum_{\substack{\nu \supset \mu\,:\,|\nu|-|\mu|=k \\ cc(\nu/\mu)=1}} (-1)^{\mathrm{ht}(\nu/\mu)}\, s_{\nu/\lambda} - \sum_{\substack{\nu \subset \lambda\,:\,|\lambda|-|\nu|=k \\ cc(\lambda/\nu)=1}} (-1)^{\mathrm{ht}(\lambda/\nu)}\, s_{\mu/\nu}\,.$$

Proof. The first statement holds more generally. Let B be a bialgebra with a countable basis $\{c_\omega\}$ containing 1, and B° its finite dual with basis $\{c^\omega\}$, such that $\langle c^\pi \mid c_\rho\rangle = \delta_{\pi\rho}$. Consider elements $f \in B^\circ$ and $g, p \in B$, such that p is primitive. Recall the canonical left action of B° on B given by $f \rightharpoonup g := \sum \langle f \mid g_2\rangle\, g_1$, where we use the Sweedler notation for the comultiplication. Similarly, there is a canonical left action of B on B°, for which the same notation is traditional. The action of symmetric functions on themselves via Hammond operators are special cases of the canonical actions of bialgebras on their duals. Since p is primitive, we have

$$c^\omega \rightharpoonup p = \begin{cases} p & \text{if } c^\omega = 1 \\ \langle c^\omega \mid p\rangle & \text{otherwise}\,. \end{cases}$$

Now recall the fact that B is a B°-module algebra under $\rightharpoonup$, and that the comultiplication can be expressed in terms of the basis $\{c_\omega\}$ as follows:

$$\delta(f) = \sum_\omega (c_\omega \rightharpoonup f) \otimes c^\omega\,.$$

Combining these facts, we have

$$\begin{aligned} f \rightharpoonup (gp) - (f \rightharpoonup g)p &= \sum_{\omega\,:\,c_\omega \neq 1} \langle c^\omega \mid p\rangle\, ((c_\omega \rightharpoonup f) \rightharpoonup g) \\ &= \left(\left(\sum_\omega \langle c^\omega \mid p\rangle\, c_\omega\right) \rightharpoonup f\right) \rightharpoonup g = (p \rightharpoonup f) \rightharpoonup g\,. \end{aligned}$$

The formula for $p_k\, s_{\mu/\lambda}$ then follows easily by letting $f = s_\lambda$, $g = s_\mu$, and by combining the previous result with (5.2) and (5.3). □

Let us now recall the principal specialization (in fact, a special case of it) of Schur functions. For every non-negative integer n, we can define a map ps^1_n from Sym_* to $\mathbb{Z}$ which sends a symmetric function $f(x_1, x_2, \ldots)$ to the integer obtained by setting $x_1 = \ldots = x_n = 1$ and $x_k = 0$ for $k > n$. In fact, we can define ps^1_n for every integer n, by letting $\mathrm{ps}^1_n(f) = \mathrm{ps}^1_{-n}(S(f))$ if $n < 0$, where S is the antipode in Sym_*. It is easy to see that ps^1_n are ring homomorphisms, and that $\mathrm{ps}^1_n(p_k) = n$ for every k. It is also known (see [19]) that

$$\mathrm{ps}^1_n(s_{\lambda'}) = \prod_{b \in \lambda} \frac{n - c(b)}{h(b)}\,. \tag{5.11}$$

We are now able to prove our result concerning the way in which $a_{\lambda\mu}$ depends on $c(\lambda, \mu)$.

Theorem 5.12. *We have*

$$a_{\lambda\mu} = \mathrm{ps}^1_{c(\lambda,\mu)} \left((\mathcal{S}^{|\pi(\lambda,\mu)|}\, s_{\pi'(\lambda,\mu)})^\perp\, s_{\rho'(\lambda,\mu)} \right)\,,$$

where $\mathcal{S}^n = \sum_{i=0}^n (-1)^i\, SQ^i$.

Proof. We let $\pi = \pi(\lambda, \mu)$, $\rho = \rho(\lambda, \mu)$, $c = c(\lambda, \mu)$, and $n = |\rho| - |\pi|$, for simplicity. We apply induction on n, which starts at $n = 1$ by Proposition 5.1. Identity (5.5) tells us that we have

$$na_{\lambda\mu} = \sum_{\lambda \subseteq \overline{\nu} \subset \mu \,:\, cc(\mu/\overline{\nu}) \leq 2} (-2)^{|\mu|-|\overline{\nu}|-1} a_{\lambda\overline{\nu}}\, d_{\overline{\nu}\mu}\,. \tag{5.13}$$

If $\mu/\overline{\nu}$ is a border strip, then $d_{\overline{\nu}\mu} = (-1)^{\mathrm{ht}(\rho/\nu)}c + d_{\nu\rho}$, where $\nu := \overline{\nu}/\pi$; otherwise $d_{\overline{\nu}\mu} = d_{\nu\rho}$. We want to lift (5.13) to an identity for symmetric functions. The crucial ingredients for our proof are the symmetric functions

$$q_k := \frac{p_k + p_{k-1} + 2p_{k-2} + \ldots + 2^{k-2}p_1}{2^{k-1}}\,,$$

which clearly satisfy $\mathrm{ps}_c^1(q_k) = c$. We claim that the following identity holds in $Sym_* \otimes \mathbb{Q}$:

$$\begin{aligned} n\,(\mathcal{S}^{|\pi|}\, s_{\pi'})^{\perp}\, s_{\rho'} = & \sum_{\substack{\pi \subseteq \nu \subset \rho \\ cc(\rho/\nu)=1}} (-2)^{|\rho|-|\nu|-1}\, \left((-1)^{\mathrm{ht}(\rho/\nu)} q_{|\rho|-|\nu|} + d_{\nu\rho}\right)\, (\mathcal{S}^{|\pi|}\, s_{\pi'})^{\perp}\, s_{\nu'} \\ & + \sum_{\substack{\pi \subseteq \nu \subset \rho \\ cc(\rho/\nu)=2}} (-2)^{|\rho|-|\nu|-1}\, d_{\nu\rho}\, (\mathcal{S}^{|\pi|}\, s_{\pi'})^{\perp}\, s_{\nu'}\,. \end{aligned} \tag{5.14}$$

If we prove this identity, the induction step is straightforward: we simply apply ps_c^1 to it, and compare the result with (5.13).

Identity (5.14) is an identity for non-homogeneous symmetric functions, so it breaks down into identities in every degree. In degree $n - i$ ($0 \leq i \leq n$), we have

$$\begin{aligned} (-1)^i n\,(SQ^i\, s_{\pi'})^{\perp}\, s_{\rho'} = & \sum_{j=0}^{i} \sum_{\substack{\nu \subset \rho \\ cc(\rho/\nu)=1}} (-1)^{j-1+|\rho|-|\nu|+\mathrm{ht}(\rho/\nu)}\, 2^{i-\alpha(j)-1}\, p_{|\rho|-|\nu|+j-i} \\ & \times (SQ^j\, s_{\pi'})^{\perp}\, s_{\nu'} + \sum_{j=0}^{i-1} \sum_{\substack{\nu \subset \rho\,:\, |\nu|=|\rho|+j-i \\ cc(\rho/\nu)\leq 2}} (-1)^{i-1}\, 2^{i-j-1}\, d_{\nu\rho}\, (SQ^j\, s_{\pi'})^{\perp}\, s_{\nu'}\,; \end{aligned} \tag{5.15}$$

here $\alpha(j)$ is j or $j-1$ depending on j being less than i or equal to i, respectively, and we adopt the convention that $p_n = 0$ if $n \leq 0$. Note that we dropped the restriction $\pi \subseteq \nu$, because if ν does not satisfy it, then $(SQ^j\, s_{\pi'})^{\perp}\, s_{\nu'} = 0$ (recall that the partitions indexing the Schur functions in the expansion of $SQ^j\, s_{\pi'}$ contain π'). For every $j = 0, \ldots, i-1$, we pair the corresponding terms in two sums in the RHS of (5.15). We start by investigating the j-th term in the second sum. We have

$$\sum_{\substack{\nu \subset \rho\,:\, |\nu|=|\rho|+j-i \\ cc(\rho/\nu)\leq 2}} d_{\nu\rho}\, s_{\nu} = D_{i-j}^{\perp}\, s_{\rho}\,.$$

We express the action of $D_{i-j}^{\perp}$ on s_ρ using Proposition 2.15 and (5.3), then we apply the standard involution on symmetric functions to the result, and obtain

$$\sum_{\substack{\nu \subset \rho\,:\, |\nu|=|\rho|+j-i \\ cc(\rho/\nu)\leq 2}} d_{\nu\rho}\, s_{\nu'} = \sum_{\substack{\nu \subset \rho \\ cc(\rho/\nu)=1}} (-1)^{j-i-1+|\rho|-|\nu|+\mathrm{ht}(\rho/\nu)}\, p_{|\rho|-|\nu|+j-i}\, s_{\nu'}\,.$$

By Lemma 5.10, when we apply the Hammond operator $(SQ^j s_{\pi'})^\perp$ to the previous expression, we can express the result as a sum of two terms, one of which is the j-th term in the first sum of the RHS of (5.15) up to a constant. Hence, by pairing the j-th terms in the two sums, we obtain

$$(-1)^{j+|\rho|}\, 2^{i-j-1} \sum_{\substack{\nu\subset\rho\\ cc(\rho/\nu)=1}} (-1)^{\mathrm{ht}(\rho/\nu)-|\nu|}\, \left(p^\perp_{|\rho|-|\nu|+j-i}\,(SQ^j\, s_{\pi'})\right)^\perp\, s_{\nu'}\,.$$

We can simplify this sum by using (5.3) again, together with the standard involution on symmetric functions:

$$\sum_{\substack{\nu\subset\rho\,:\,|\nu|=k\\ cc(\rho/\nu)=1}} (-1)^{\mathrm{ht}(\rho/\nu)} s_{\nu'} = (-1)^{|\rho|-k-1}\, p^\perp_{|\rho|-k}\, s_{\rho'}\,. \tag{5.16}$$

Hence, by the commutativity of Hammond operators, the sum of the j-th terms in the two sums in the RHS of (5.15) can be expressed as

$$(-1)^{j-1}\, 2^{i-j-1} \left(\left(\sum_k p_{|\rho|-k}\, p^\perp_{|\rho|-k+j-i}\right)\,(SQ^j\, s_{\pi'})\right)^\perp s_{\rho'}\,; \tag{5.17}$$

here the summation ranges over all k with $|\rho|-k$ lying between 1 and the length of the longest border strip of ρ. We want to show that we can actually extend the range of $|\rho|-k$ to infinity without affecting the result; this has the advantage of being able to write the sum as the operator D_{i-j}. We regard $SQ^j\, s_{\pi'}$ as a linear combination of Schur functions s_σ with $|\sigma| = |\pi| + j$ and $\pi' \subseteq \sigma$. Applying D_{i-j} viewed as $\sum_{l\geq 1} p_l\, p^\perp_{l+j-i}$ to s_σ gives a sum of Schur functions s_τ, where τ is obtained from σ by removing a border strip of length $l+j-i>0$, and then adding a border strip of length l. Since we then consider $s_{\rho'/\tau}$, we are only interested in those τ contained in ρ'. Clearly, for such τ, the corresponding added border strip has length l at most the length of the longest border strip of ρ, that is, within the range of $|\rho|-k$. According to the above remarks and (5.5), we can express the RHS of (5.15) minus the term corresponding to $j=i$ in the first sum as

$$\sum_{j=0}^{i-1} (-1)^{j-1}\, 2^{i-j-1}\, ((D_{i-j}\, SQ^j)\, s_{\pi'})^\perp\, s_{\rho'} = (-1)^i\, i\, (SQ^i\, s_{\pi'})^\perp\, s_{\rho'}\,. \tag{5.18}$$

It only remains to deal with the term corresponding to $j=i$ in the RHS of (5.15). Applying (5.15) again, the commutativity of Hammond operators, and Lemma 5.9, we can express this term as follows:

$$(-1)^i \sum_{\substack{\nu\subset\rho\\ cc(\rho/\nu)=1}} (-1)^{|\rho|-|\nu|-1+\mathrm{ht}(\rho/\nu)}\, p_{|\rho|-|\nu|}\, (SQ^i\, s_{\pi'})^\perp\, s_{\nu'} \tag{5.19}$$

$$= (-1)^i \left(\sum_k p_{|\rho|-k}\, p^\perp_{|\rho|-k}\right) \left((SQ^i\, s_{\pi'})^\perp\, s_{\rho'}\right) = (-1)^i (n-i)\, (SQ^i\, s_{\pi'})^\perp\, s_{\rho'}\,;$$

here the range of $|\rho|-k$ is the same as in (5.17), and was extended to infinity by the same argument as above. Combining (5.18) and (5.19) proves (5.15). □

Theorem 5.12 says that $a_{\lambda\mu}$ is a certain specialization of a non-homogeneous symmetric function with highest homogeneous component $s_{\mu'/\lambda'}$. More precisely, $a_{\lambda\mu}$ is a polynomial in $c(\lambda,\mu)$ depending only on μ/λ; it has degree $|\mu|-|\lambda|$, free term equal to $a_{\pi(\lambda,\mu)\rho(\lambda,\mu)}$, and coefficient of the leading term equal to the corresponding coefficient of $\mathrm{ps}^1_{c(\lambda,\mu)}(s_{\mu'/\lambda'})$. Thus, we have reduced the computation of $a_{\lambda\mu}$ to the case $l(\lambda)<l(\mu)$, $\lambda_1<\mu_1$. One can use Proposition 5.6 to compute $SQ^i\, s_{\pi'(\lambda,\mu)}$ for all $i=1,\ldots,|\pi(\lambda,\mu)|$ (in fact we only need the coefficients $a_{\pi'(\lambda,\mu)\nu}$ with $\nu\subseteq\rho'(\lambda,\mu)$). Then Theorem 5.12 enables us to compute quite easily the coefficient $a_{\lambda\mu}$. Of course, the complexity is in the previous step, and it increases with the weight of $\pi(\lambda,\mu)$. The following Corollary considers the cases when this weight is 0 or 1.

Corollary 5.20. *If $\nu:=\mu/\lambda$ is a Young diagram, then $a_{\lambda\mu}=\mathrm{ps}^1_{c(\lambda,\mu)}(s_{\nu'})$, which can be expressed using (5.11). If $\pi(\lambda,\mu)$ consists of a single box, then $a_{\lambda\mu}=\mathrm{ps}^1_{c(\lambda,\mu)}((p_1-p_2)^{\perp}\, s_{\rho'(\lambda,\mu)})$.*

Proof. The above statements are immediate consequences of Theorem 5.12. All we need to observe (for the second part) is that $SQ^1\, s_1=D^1\, p_1=p_2$, by Proposition 2.15. □

The following Corollary gives sufficient conditions for a coefficient $a_{\lambda\mu}$ to be 0. To state it, we need additional notation. For every partition λ and non-negative integer n, we denote by $m(\lambda,n)$ the largest number k for which either $\lambda'_k+\ldots+\lambda'_{l(\lambda)}>n$ or $\lambda'_{k+1}+\ldots+\lambda'_{l(\lambda)}=n$; if no such k exists, we let $m(\lambda,n):=0$.

Corollary 5.21. *Given diagrams $\lambda\subset\mu$, we consider the partitions σ and τ whose parts are the lengths of columns and rows of μ/λ, respectively. The coefficient $a_{\lambda\mu}$ is 0 whenever the following condition is satisfied:*

$$1-m(\sigma,|\pi(\lambda,\mu)|)\leq c(\lambda,\mu)\leq m(\tau,|\pi(\lambda,\mu)|)-1\,.$$

Proof. We let $\pi=\pi(\lambda,\mu)$ and $\rho=\rho(\lambda,\mu)$, for simplicity. Clearly, it suffices to consider the case $2|\pi|<|\rho|$, since otherwise $m(\sigma,|\pi|)=m(\theta,|\pi|)=0$, so the above condition cannot hold. Given i between 0 and $|\pi|$, we view $(SQ^i\, s_{\pi'})^{\perp}\, s_{\rho'}$ as a linear combination of skew Schur functions $s_{\rho'/\nu}$, where $\pi'\subseteq\nu\subseteq\rho'$ and $|\nu|-|\pi|=i$. We then view every $s_{\rho'/\nu}$ as a linear combination (with non-zero coefficients) of Schur functions $s_{\theta'}$. By the Littlewood-Richardson rule, the minimum value $m_c(\nu)$ of θ_1 for a fixed ν is equal to the length of the longest column of the skew shape ρ'/ν. Indeed, this is clearly a lower bound, since the boxes in the mentioned column have to be filled with different symbols in any Littlewood-Richardson filling; on the other hand, the lower bound is attained, since by filling every column of ρ'/ν with symbols $1,2,\ldots$ (from top to bottom, in this order), we obtain a Littlewood-Richardson filling. By applying the standard involution on symmetric functions, we can deduce from here that the minimum value $m_r(\nu)$ of $l(\theta)$ for a fixed ν is equal to the length of the longest row of the skew shape ρ'/ν. By (5.11), we have that $\mathrm{ps}^1_{c(\lambda,\mu)}(s_{\rho'/\nu})=0$ whenever $1-m_r(\nu)\leq c(\lambda,\mu)\leq m_c(\nu)-1$. To minimize $m_r(\nu)$, we let $i=|\pi|$, and construct a sequence of Young diagrams $\pi'=\nu_{(0)},\nu_{(1)},\ldots,\nu_{(|\pi|)}$ such that every $\nu_{(j)}$ is obtained by adding a box to one of the rows of $\rho'/\nu_{(j-1)}$ of maximum length. It is not difficult to see that there is always such a choice, and that $m_r(\nu_{(|\pi|)})=m(\sigma,|\pi|)$ minimizes $m_r(\nu)$. The same argument applies to $m_c(\nu)$, thus proving the Corollary. □



Note that if $|\mu| - |\lambda| \leq 2$, the condition in the Corollary is also necessary (which is not true in general, however). Indeed, if $|\mu| - |\lambda| = 1$, the necessity is clear. If $|\mu| - |\lambda| = 2$, there are three cases possible: (1) $\mu/\lambda = (2)$, (2) $\mu/\lambda = (1,1)$, and (3) μ/λ is a skew diagram with two boxes of content c_1, c_2. By Proposition 5.6, we have $a_{\lambda\mu} = c(\lambda,\mu)\,(c(\lambda,\mu)-1)/2$ in the first case, $a_{\lambda\mu} = c(\lambda,\mu)\,(c(\lambda,\mu)+1)/2$ in the second one, and $a_{\lambda\mu} = c_1c_2 + 1$ in the third one. The necessity of the condition in the Corollary follows without difficulty from here.

Example 5.22. Let $\pi = (3,2,2,1)$, $\rho = (9,7,6,5,4,4,2)$, and λ, μ be such that $\pi(\lambda,\mu) = \pi$ and $\rho(\lambda,\mu) = \rho$. We have $\sigma = (6,5,4,4,3,3,2,1,1)$ and $\tau = (6,5,4,4,4,4,2)$, whence $m(\sigma,8) = 3$ and $m(\tau,8) = 4$. According to the Corollary, we have $a_{\lambda\mu} = 0$ for all λ, μ with $-2 \leq c(\lambda,\mu) \leq 3$. In the figure above, illustrating this example, we marked with a dot the boxes where the top left box of ρ can be translated, such that the corresponding coefficient $a_{\lambda\mu}$ is 0.

The next logical step after Theorem 5.12 would be to investigate how $a_{\lambda\mu}$ depends on the contents $c_i(\lambda,\mu)$ corresponding to the connected components of μ/λ (assuming that this is not connected). It is known that if μ/λ consists of at least two connected components (each of which is a skew diagram), then $s_{\mu/\lambda}$ is the product of the corresponding skew Schur functions. This property does not seem to have a straightforward analogue for the coefficients $a_{\lambda\mu}$, which not only depend on μ/λ, but also on $c(\lambda,\mu)$. We conclude this section with a conjecture which represents such an analogue when μ/λ has two connected components. Before stating it, we introduce the following notation involving two Young diagrams λ, μ, and two integers k, n:

$$f(\lambda,\mu,k,n) := \sum_{\nu\subseteq\lambda,\,\nu\subseteq\mu} \left(\prod_{b\in\lambda/\nu} \frac{k-c(b)}{h(b)}\right)\left(\prod_{b\in\mu/\nu} \frac{n-c(b)}{h(b)}\right);$$

here ν can be the empty diagram, and if $\nu = \lambda$ (or $\nu = \mu$), the corresponding product is considered to be 1.

Conjecture 5.23. *Let $\lambda \subset \mu$ be such that μ/λ is a union of two connected components. Let*

$$\left(\mathcal{S}^{|\pi_{(k)}(\lambda,\mu)|}\, s_{\pi'_{(k)}(\lambda,\mu)}\right)^{\perp} s_{\rho'_{(k)}(\lambda,\mu)} = \sum_{\nu \in P_k} \alpha_{k,\nu}\, s_{\nu'}\,,$$

for $k = 1, 2$, where P_k are corresponding sets of partitions. Then we have

$$a_{\lambda\mu} = \sum_{\substack{\nu_{(1)} \in P_1 \\ \nu_{(2)} \in P_2}} \alpha_{1,\nu_{(1)}}\, \alpha_{2,\nu_{(2)}}\, f(\nu_{(1)}, \nu_{(2)}, c_1(\lambda,\mu), c_2(\lambda,\mu))\,.$$

This conjecture was tested for small diagrams $\pi_{(k)}, \rho_{(k)}$, $k = 1, 2$. There are indications that similar formulas exist for three or more connected components. Such formulas would reduce the computation of $a_{\lambda\mu}$ to the case $l(\lambda) < l(\mu)$, $\lambda_1 < \mu_1$, and μ/λ having only one connected component, whence they would increase the efficiency of the computation with respect to the formula in Theorem 5.12.

6. Applications to the Geometry of Grassmannians

Corollary 5.20 provides a closed formula for the action of Steenrod operations on Schubert classes of the form $\sigma_{(j^{n-i}k^i)}$ and $\sigma_{(j^{n-i-1}(j+1)k^i)}$ in the cohomology of $Gr_n(\mathbb{C}^{n+k})$. Indeed, since Schubert classes in the cohomology of this Grassmannian are indexed by partitions with at most n rows and k columns, all the coefficients $a_{\lambda\mu}$ we need to compute are of the form specified by the Corollary. In general, we can combine the above results (Proposition 5.6, Theorem 5.12, and possibly Conjecture 5.23) to compute the action of Steenrod operations on any Schubert class.

Let us consider some examples. Since the cell structure and the action of the Steenrod algebra on projective spaces are well-known, we consider the Grassmannian $Gr_2(\mathbb{C}^5)$ of dimension 12, and the Schubert cells

$$\Omega_{(1,1)} := \{V \subset \mathbb{C}^3 \,:\, \dim(V \cap \mathbb{C}^2) \geq 1\}\,, \quad \Omega_{(3,1)} := \{V \,:\, \dim(V \cap \mathbb{C}^2) \geq 1\}$$

of dimensions 4 and 8, respectively. By Corollary 5.20, we have

$$Sq^4\, \sigma_{(3,1)} = 0\,, \quad (Sq^2)^2\, \sigma_{(3,1)} = 0\,, \quad \mathcal{P}^1\, \sigma_{(3,1)} = 0 \ \text{(for } p = 3).$$

The integral lifts of these operations also give 0. This suggests that the top cell $\Omega_{(3,3)}$ of $Gr_2(\mathbb{C}^5)$ (the space itself) might be attached trivially to $\Omega_{(3,1)}$ (cf. the discussion in §3). On the other hand, we have

$$Sq^4\, \sigma_{(1,1)} = \sigma_{(2,2)}\,, \quad (Sq^2)^2\, \sigma_{(1,1)} = 0\,;$$

in fact, the integral lift gives $(SQ^1 \circ SQ^1)\, s_{(1,1)} = 2 s_{(3,1)}$. However,

$$\mathcal{P}^1\, \sigma_{(1,1)} = \sigma_{(3,1)} - \sigma_{(2,2)} \ \text{(for } p = 3),$$

which means that $\Omega_{(3,1)}$ is attached non-trivially to $\Omega_{(1,1)}$.

The previous example confirms that the mod 2 Steenrod operations give only partial information about the attaching maps. However, they provide a good first approximation of the cell structure of a space (see below). Then, one has to use more sophisticated

methods to decide whether a cell which does not appear in any composition of Steenrod squares on a lower dimensional cell is attached trivially to the latter or not. For instance, one might start by considering Steenrod operations corresponding to primes $p > 2$, or Adams operations in K-theory etc. It is also important to distinguish between unstable and stable attachment. It would be interesting to see whether the cell Ω_μ is attached trivially to Ω_λ for λ and μ satisfying the condition in Corollary 5.21; it might be possible that it is, at least in some cases, as the fact that $a_{\lambda\mu} = 0$ (integrally!) suggests. A more difficult problem is to determine the elements in the homotopy groups of spheres corresponding to the *relative attaching maps*; it is known that for projective spaces they are the images of the so-called *J-homomorphism*.

The approximation of the cell structure of a space provided by the Steenrod squares can be represented by a directed graph with weighted edges; we consider the cells as vertices and an edge of weight $2(|\mu| - |\lambda|)$ from Ω_λ to Ω_μ if and only if $a_{\lambda\mu} = 1 \bmod 2$.

Remarks 6.1.

(1) *The coefficient of σ_μ in $Sq^{i_1} \dots Sq^{i_k} \sigma_\lambda$ is equal (mod 2) to the number of paths with edge weights $i_1, \dots, i_k$ (in this order) from Ω_λ to Ω_μ in the directed graph mentioned above. Hence, we can recover the action of every composition of Steenrod squares from the graph.*
(2) *The well-known fact that the Steenrod squares Sq^{2^k}, $k \geq 0$, generate the mod 2 Steenrod algebra implies that we do not lose information by removing all edges of the graph whose weights are not powers of 2.*

Proceeding as shown above, we can easily draw the directed graph (in Remark 6.1 (2)) corresponding to the Grassmannian $Gr_2(\mathbb{C}^5)$ (see the figure below); the cells are represented by the Young diagrams corresponding to the partitions which index them, and the edge weights are obvious. Note that we obtained $Sq^4 \sigma_{(2)} = 0$ by Corollary 5.6; all the other computations are based on the simpler expressions provided by Corollary 5.20.



We conclude with one more remark, related to the so-called *Steenrod length* of Grassmannians. The mod 2 Steenrod length of an object X in the stable homotopy category was defined in [8] as one more than the length of the longest chain of non-zero Steenrod operations in the mod 2 cohomology of X. For instance, the Steenrod length of $Gr_2(\mathbb{C}^5)$ is 4 because $Sq^2Sq^4Sq^2\,\sigma_{(2)} = \sigma_{(3,3)}$ and $Sq^2Sq^4(Sq^2)^2\,\sigma_{(1)} = 0$. The Steenrod length of X is no greater than the so-called *ghost length* of X, which is a more accurate measure of the complexity of X, but more difficult to compute (see [8]). The Steenrod length of real projective spaces was computed by Vakil in [32]. We suggest that our formulas could help extend his work to arbitrary Grassmannians. However, the general case will be considerably harder, at least for two reasons: (1) we need mod 2 congruences for the coefficients $a_{\lambda\mu}$, for which the only expression we have at this time is the one in Corollary 5.6 if μ/λ is skew; (2) in general, there is more than one path with prescribed edge weights between two vertices in the directed graph discussed above, as opposed to at most one path in the case of projective spaces.

References

[1] J. F. Adams. *Stable Homotopy and Generalised Homology*. Chicago University Press, Chicago, 1972.

[2] J. M. Boardman. The eightfold way to BP-operations. In *Current Trends in Algebraic Topology*, volume 2 of *Canad. Math. Soc. Conf. Proc.*, pages 187–226, Providence, RI, 1982. Amer. Math. Soc.

[3] A. Borel and J.-P. Serre. Groupes de Lie et puissances réduites de Steenrod. *Amer. J. Math.*, 75:409–448, 1953.

[4] E. Brown, D. Davis, and F. Peterson. The homology of BO and some results about the Steenrod algebra. *Math. Proc. Cambridge Philos. Soc.*, 81:393–398, 1977.

[5] E. Brown and F. Peterson. Some remarks about symmetric functions. *Proc. Amer. Math. Soc.*, 60:349–352, 1976.

[6] D. P. Carlisle and G. Walker. Poincaré series for the occurence of certain modular representations of $GL(n, p)$ in the symmetric algebra. *Proc. Royal Soc. Edinburgh*, 113A:27–41, 1989.

[7] L. Carlitz. Some inversion formulas. *Rend. Circ. Mat. Palermo*, 12:183–199, 1963.
[8] J. D. Christensen. *Ideals in triangulated categories: Phantoms, ghosts and skeleta.* PhD thesis, M.I.T., 1997.
[9] P. Doubilet. On the foundations of combinatorial theory VII: Symmetric functions through the theory of distribution and occupancy. *Studies Appl. Math.*, LI:377–396, 1972.
[10] W. Fulton. *Young Tableaux*, volume 35 of *London Math. Soc. Student Texts*. Cambridge Univ. Press, Cambridge and New York, 1997.
[11] H. Hiller. *Geometry of Coxeter Groups*, volume 54 of *Res. Notes in Math.* Pitman Books Ltd., London, 1982.
[12] V. G. Kac and A. K. Raina. *Bombay Lectures on Highest Weight Representations of Infinite Dimensional Lie Algebras*, volume 2 of *Adv. Ser. in Math. Phys.* World Scientific, Singapore and New Jersey, 1987.
[13] M. Kaneda, M. Shimada, M. Tezuka, and N. Yagita. Representations of the Steenrod algebra. *J. Algebra*, 155:435–454, 1993.
[14] T. Katsura, Y. Shimizu, and K. Ueno. Complex cobordism ring and conformal field theory over $\mathbb{Z}$. *Math. Ann.*, 291:551–571, 1991.
[15] N. J. Kuhn. Generic representations of the finite general linear groups and the Steenrod algebra: I. *Amer. J. Math.*, 116:327–360, 1993.
[16] A. Kulikauskas and J. B. Remmel. Primitive bi-brick permutations and symmetric functions. Preprint, 1994.
[17] T. Lance. Steenrod and Dyer-Lashof operations on BU. *Trans. Amer. Math. Soc.*, 276:497–510, 1983.
[18] P. S. Landweber. Cobordism operations and Hopf algebras. *Trans. Amer. Math. Soc.*, 27:94–110, 1967.
[19] I. G. Macdonald. *Symmetric Functions and Hall Polynomials*. Oxford Mathematical Monographs. Oxford University Press, Oxford, second edition, 1995.
[20] P. A. MacMahon. *Combinatory Analysis*. Chelsea, New York, reprint edition, 1960.
[21] J. Milnor. The Steenrod algebra and its dual. *Ann. Math.*, 67:150–171, 1958.
[22] J. Milnor. On the cobordism ring Ω_* and a complex analogue. *Amer. J. Math.*, 82:505–521, 1960.
[23] S. Montgomery. *Hopf Algebras and Their Action on Rings*, volume 82 of *Regional Conf. Ser. in Math.* Amer. Math. Soc., Providence, RI, 1993.
[24] S. P. Novikov. The methods of algebraic topology from the point of view of cobordism theory. *Izv. Akad. Nauk SSSR, Ser. Mat.*, 31:855–951, 1967.
[25] F. Peterson. A mod p Wu formula. *Bol. Soc. Mat. Mexicana*, 20:56–58, 1975.
[26] N. Ray. Private communication, February 1995.
[27] W. R. Schmitt. Incidence Hopf algebras. *J. Pure Appl. Algebra*, 96:299–330, 1994.
[28] L. Schwartz. *Unstable Modules over the Steenrod Algebra and Sullivan's Fixed Point Conjecture*. Chicago Lecture Notes in Math. Univ. of Chicago Press, Chicago and London, 1994.
[29] B. Shay. Mod p Wu formulas for the Steenrod algebra and the Dyer-Lashof algebra. *Proc. Amer. Math. Soc.*, 63:339–347, 1977.
[30] N. E. Steenrod and D. B. A. Epstein. *Cohomology Operations*. Ann. of Math. Stud. Princeton Univ. Press, Princeton, NJ, 1962.
[31] T. Sugawara. Wu formulas for the mod 3 reduced power operations. *Mem. Fac. Sci. Kyushu Univ. Ser. A*, 33:297–309, 1979.
[32] R. Vakil. On the Steenrod length of real projective spaces: finding longest chains in certain directed graphs. Preprint, Princeton University, 1997.
[33] R. M. W. Wood. Differential operators and the Steenrod algebra. Preprint, University of Manchester, 1996.
[34] W. T. Wu. Les i-carrés dans une varieté grassmannième. *C. R. Acad. Sci. Paris*, 230:918–920, 1950.

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139

E-mail address: lenart@math.mit.edu