

# Nucleation rate of particles in the lower atmosphere: Estimated time needed to reach pseudo-steady state and sensitivity to H<sub>2</sub>SO<sub>4</sub> gas concentration

Fangqun Yu

Atmospheric Sciences Research Center, State University of New York, Albany, New York, USA

Received 8 February 2003; revised 10 April 2003; accepted 16 April 2003; published 24 May 2003.

[1] Using a kinetic model, we estimate the time-needed ( $t_0$ ) to establish a pseudo-steady state nucleation when sulfuric acid concentration ([H<sub>2</sub>SO<sub>4</sub>]) suddenly increases to a certain fixed level. We find that  $t_0$  is roughly inversely proportional to [H<sub>2</sub>SO<sub>4</sub>]. The assumption of instantaneously achieved pseudo-steady state concentration of critical clusters, on which the classical nucleation theories are based, is no longer valid when  $t_0$  is longer than the time period that [H<sub>2</sub>SO<sub>4</sub>] can be treated as constant. Under the lower marine atmospheric conditions considered in this study, the nucleation rate predicted by ion-mediated nucleation theory is very sensitive to [H<sub>2</sub>SO<sub>4</sub>] when [H<sub>2</sub>SO<sub>4</sub>] is low ( $< \sim 7 \times 10^6/\text{cm}^3$ ) but such a sensitivity reduces significantly as [H<sub>2</sub>SO<sub>4</sub>] increases, which is quite different from that predicted with classical binary and ternary homogeneous nucleation theories. **INDEX TERMS:** 0305 Atmospheric Composition and Structure: Aerosols and particles (0345, 4801); 0365 Atmospheric Composition and Structure: Troposphere—composition and chemistry. **Citation:** Yu, F., Nucleation rate of particles in the lower atmosphere: Estimated time needed to reach pseudo-steady state and sensitivity to H<sub>2</sub>SO<sub>4</sub> gas concentration, *Geophys. Res. Lett.*, 30(10), 1526, doi:10.1029/2003GL017084, 2003.

## 1. Introduction

[2] Despite intensive research over last several decades, the fundamental mechanism that leads to the ubiquitous background of ultrafine aerosols in the atmosphere remains poorly understood. Binary homogeneous nucleation (BHN) of H<sub>2</sub>SO<sub>4</sub> and H<sub>2</sub>O has generally been presumed to be the principal particle formation mechanism. The difficulty of BHN theory in explaining the particle formation observed recently in the lower troposphere implies that mechanisms other than BHN may be responsible for the observed nucleation events [e.g., *Weber et al.*, 1996; *Clarke et al.*, 1998]. Ternary homogeneous nucleation (THN) theory involving H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O-NH<sub>3</sub> [e.g., *Coffman and Hegg*, 1995; *Korhonen et al.*, 1999; *Ball et al.*, 1999; *Napari et al.*, 2002] and ion-mediated nucleation (IMN) involving H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O-Ion [*Yu and Turco*, 2000] have been proposed as the alternative particle formation mechanisms.

[3] Based on THN theory, ammonia stabilizes the critical embryo (reducing its size) and thus dramatically increases the nucleation rate. According to IMN theory, the charged molecular clusters, formed around air ions, are much more stable and can grow significantly faster than corresponding

neutral clusters, and thus can preferentially achieve stable, observable sizes [*Yu and Turco*, 2000, 2001]. One of major differences between IMN theory and classical nucleation theories is that, at a given sulfuric acid gas concentration ([H<sub>2</sub>SO<sub>4</sub>]), IMN theory considers kinetically how clusters of various sizes evolve starting at monomers while classical theories assumes an instantaneous pseudo-steady state concentration of critical embryos. [H<sub>2</sub>SO<sub>4</sub>] usually changes rapidly during the nucleation events, and question concerning the validity of instantaneous pseudo-steady state assumption has been raised [*Yu and Turco*, 2001].

[4] While both THN and IMN theories introduce new specie into the nucleation system (NH<sub>3</sub> and ions, respectively), H<sub>2</sub>SO<sub>4</sub> still is the key nucleating specie. In this paper, we study the sensitivity of nucleation rate to [H<sub>2</sub>SO<sub>4</sub>]. The time-needed to establish the pseudo-steady state nucleation when [H<sub>2</sub>SO<sub>4</sub>] suddenly increases to a certain level is also investigated. Our emphasis is on IMN theory but comparisons with BHN and THN theories are made.

## 2. IMN Theory and Kinetic Model

[5] The particle nucleation is essentially the result of several competing dynamic processes of molecular clusters: clustering growth, evaporation shrink, and scavenging by pre-existing particles. In the atmosphere, the clustering among hydrated H<sub>2</sub>SO<sub>4</sub> gas molecules leads to the formation and growth of various clusters. Those clusters that survive the evaporation/scavenging and grow to stable sizes are considered nucleated particles. Under typical conditions in the lower atmosphere where nucleation events have been observed, neutral sub-critical H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O clusters are very unstable and evaporate quickly back to monomers.

[6] Both H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O-NH<sub>3</sub> and H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O-Ion clusters are believed to be much more stable than neutral H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O clusters. In addition, H<sub>2</sub>SO<sub>4</sub>-H<sub>2</sub>O-Ion clusters grow much faster than the neutral ones as a result of electrical interactions of polar molecules with ions [*Yu and Turco*, 2000, 2001]. The charged clusters lose these advantages (enhanced stability and growth rate) when they are neutralized by ions of opposite sign. The dynamic interactions among gas molecules and clusters of both charged and neutral are the essentials of IMN theory and only a kinetic model can simulate these processes properly [*Yu and Turco*, 2001].

[7] We employ an advanced particle microphysics model that simulates a size-resolved multicomponent aerosol system via a unified collisional mechanism involving both neutral and charged particles down to molecular sizes [*Yu*

and Turco, 2001]. The size-resolved ion-ion recombination coefficients, ion-neutral collision kernels, and neutral-neutral interaction coefficients calculated in the model are physically consistent and depend on ambient conditions [Yu and Turco, 2001]. Due to the lack of the information about the evaporation coefficients of small clusters, a simple adjustment to the accommodation coefficient was assumed as an expedient means of parameterizing the evaporation process in our earlier model [Yu and Turco, 2001]. In the current model, the evaporation equations [Yu and Turco, 2001] are solved explicitly, with the evaporation coefficients of neutral  $\text{H}_2\text{SO}_4\text{-H}_2\text{O}$  clusters calculated based on the assumption that the escape frequency of  $\text{H}_2\text{SO}_4$  molecules equals the collision frequency when the cluster is in equilibrium with the surrounding  $\text{H}_2\text{SO}_4$  molecules. Nevertheless, for the marine boundary conditions studied here, two different approaches (to treat the evaporation process) give similar results as both approaches indicate that the growth of neutral sub-critical clusters is too slow to lead to any nucleation. The scheme to calculate enhancement factors for the interactions rates of  $\text{H}_2\text{SO}_4$  molecules with ions/charged particles has been updated with a recently derived analytical expression which gives the enhancement factor as a function of the size of the charged particles, the size and dipole moment of condensable vapor molecules, and ambient temperature [Nadykto and Yu, 2003].

[8] It is noteworthy that IMN theory is different from the classical ion-nucleation (INU) theory [e.g., Hamill et al., 1982; Raes et al., 1986]. Classical “ion-nucleation” modifies the free energy needed to form critical embryos (similar to THN) by considering the Thomson effect, but does not consider how the pseudo-steady state distribution of clusters is established. Most importantly, the effect of charge on the growth rate of clusters, which is critical, is not considered.

[9] At a given condition, the classical nucleation theories (BHN, THN, INU) predict the nucleation rate from formulas assuming a pseudo-steady state concentration of critical embryos. On the contrast, the kinetic model on which the IMN theory is based explicitly simulates the evolution of cluster distributions. From these varying cluster distributions ( $dN/dD_p$ ), an equivalent IMN theory-predicted nucleation rate ( $J_{D_0}^{\text{IMN}}$ ) of particles with diameter  $D_0$  can be calculated as the following,

$$J_{D_0}^{\text{IMN}} = \left. \frac{dN}{dD_p} \right|_{D_0} \frac{dD_p}{dt} \Big|_{D_0} = \left. \frac{dN}{dD_p} \right|_{D_0} \cdot 2v_1 \cdot \frac{[\text{H}_2\text{SO}_4]}{4} \cdot \bar{c} \quad (1)$$

where  $\bar{c}$  is the mean thermal speed of the condensing hydrated  $\text{H}_2\text{SO}_4$  monomers, and  $v_1$  is the amount that particle volume increases upon the addition of a single  $\text{H}_2\text{SO}_4$  molecule (and associated  $\text{H}_2\text{O}$  molecules to maintain phase equilibrium). Two values of  $D_0$  are considered here: diameter of critical embryo ( $D_{\text{crit}}$ ) and 3 nm. Equation (1) is the same as the formula Weber et al. [1996] used to estimate observed nucleation rates ( $J_{3\text{nm}}^{\text{obs}}$ ) from the measured  $[\text{H}_2\text{SO}_4]$  and particle size distributions down to 3 nm.

### 3. Simulations

[10] We consider two different ambient conditions corresponding to two different nucleation events observed: one in the tropical marine boundary layer [Clarke et al., 1998, — Case1] and the other at a remote marine site [Mauna Loa

Observatory, Weber et al., 1996, — Case 2]. The mean values of temperature ( $T$ ), relative humidity ( $RH$ ), and pre-existing particle surface area ( $S_0$ ) are  $25^\circ\text{C}$ , 95%,  $10 \mu\text{m}^2/\text{cm}^3$ , respectively for case 1 and are  $8^\circ\text{C}$ , 48%,  $50 \mu\text{m}^2/\text{cm}^3$ , respectively for case 2. Based on the locations of the measurement sites, we choose an ionization rate ( $Q$ ) of 2 and 5 ion-pairs/ $\text{cm}^3\text{s}$  [Reiter, 1992] for case 1 and case 2, respectively. Two additional ionization rates are considered for sensitivity study (5 and 10 ion-pairs/ $\text{cm}^3\text{s}$  for case 1; 2 and 10 ion-pairs/ $\text{cm}^3\text{s}$  for case 2).

[11] Three types of particles (neutral, positively charged, and negatively charged) are treated in the model. All particles are assumed to be composed of  $\text{H}_2\text{SO}_4$  and  $\text{H}_2\text{O}$  and the water contents of clusters/particles are calculated based on thermodynamic equilibrium. We initialized our model with neutral monomers (i.e, hydrated  $\text{H}_2\text{SO}_4$  gas molecules) and the ambient particles corresponding to the surface area given above. At  $t = 0$  minute, only monomers and ambient particles exist. As time passes by, monomers aggregate into clusters of various sizes. Our model keeps track the evolution (growth, neutralization, evaporation, scavenging, etc) of the size distributions of both charged and neutral clusters/particles. From these size distributions, the nucleation rates are calculated with equation (1).


#### 3.1. Estimated Time Needed to Achieve Pseudo-Steady State

[12] Figure 1 shows the variations of  $J_{D_{\text{crit}}}^{\text{IMN}}$  (solid lines) and  $J_{3\text{nm}}^{\text{IMN}}$  (dot-dashed lines) with time at 4 different values of  $[\text{H}_2\text{SO}_4]$ : (a) case 1, and (b) case 2. We run the model for three hours and the value of  $[\text{H}_2\text{SO}_4]$  for each run is indicated along each line.  $T$ ,  $RH$ ,  $Q$ , and  $[\text{H}_2\text{SO}_4]$  are fixed during each 3-hour simulation. The diameter of critical embryo ( $D_{\text{crit}}$ , calculated with classical nucleation theory) increases as  $[\text{H}_2\text{SO}_4]$  decreases. The values of  $J_{D_{\text{crit}}}^{\text{IMN}}$  are calculated at a specific  $D_{\text{crit}}$  corresponding to each  $[\text{H}_2\text{SO}_4]$ . As expected,  $J = 0/\text{cm}^3\text{s}$  at the beginning since no clusters of stable sizes have formed yet. The nucleation rates increase quickly as clusters begin to form and reach relative stable values (i.e, pseudo-steady state) after a certain period of time ( $t_0$ ).  $t_0$  can be defined as the time when the relative change of nucleation rate becomes smaller than certain value  $\varepsilon_0$  for the first time,

$$\left. \frac{1}{J_D^{\text{IMN}}} \frac{dJ_D^{\text{IMN}}}{dt} \right|_{t_0} < \varepsilon_0 \quad (2)$$

[13] We choose  $\varepsilon_0 = 0.6\% \text{ min}^{-1}$ , and the position of  $t_0$  in each line is also indicated in Figure 1 (filled circles for  $J_{D_{\text{crit}}}^{\text{IMN}}$  and opaque circles for  $J_{3\text{nm}}^{\text{IMN}}$ ). An  $\varepsilon_0$  value of 0.6% gives visually reasonable locations of  $t_0$  in the curves shown in Figure 1 that separate the regions with rapid change in nucleation rate and regions with stable nucleation rate.

[14] The nucleation system achieves pseudo-steady state (with regard to formation rate of particles with diameters  $D_{\text{crit}}$  and 3nm) quickly when  $[\text{H}_2\text{SO}_4]$  is high. For the two cases considered in this study,  $t_0$  for  $J_{D_{\text{crit}}}^{\text{IMN}}$  is  $\leq 5$  minutes and for  $J_{3\text{nm}}^{\text{IMN}}$  is  $\leq 15$  minutes at  $[\text{H}_2\text{SO}_4]$  of  $5 \times 10^7/\text{cm}^3$ . Note that the slight decrease in nucleation rates after  $t_0$  at  $[\text{H}_2\text{SO}_4]$  of  $5 \times 10^7/\text{cm}^3$  is due to the scavenging of sub-nucleated clusters by nucleated particles.  $t_0$  is almost inversely proportional to  $[\text{H}_2\text{SO}_4]$ . When  $[\text{H}_2\text{SO}_4] = 1 \times$



**Figure 1.** The variations of model calculated  $J_{D_{crit}}^{IMN}$  (solid lines) and  $J_{3nm}^{IMN}$  (dot-dashed lines) with time at 4 different values of  $[H_2SO_4]$ : (a) case 1 ( $T=25^\circ C$ ,  $RH=95\%$ ,  $S_0=10 \mu m^2/cm^3$ ,  $Q=2$  ion-pairs/ $cm^3 s$ ), and (b) case 2 ( $T=8^\circ C$ ,  $RH=48\%$ ,  $S_0=50 \mu m^2/cm^3$ ,  $Q=5$  ion-pairs/ $cm^3 s$ ). The value of  $[H_2SO_4]$  for each run is indicated along each line.  $T$ ,  $RH$ ,  $Q$ , and  $[H_2SO_4]$  are fixed during each 3-hour simulation. The corresponding values of time needed to established pseudo-steady state nucleation are also indicated.

$10^7/cm^3$ ,  $t_0$  is  $\sim$  half hour for  $J_{D_{crit}}^{IMN}$  and  $\sim$  one hour for  $J_{3nm}^{IMN}$ .


[15] In our simulations, we assume that  $[H_2SO_4]$  is constant. In the real atmosphere where significant nucleation has been observed,  $[H_2SO_4]$  typically increases rapidly and then decreases quickly [e.g., *Weber et al.*, 1996; *Clarke et al.*, 1998]. There exist a phase lag between the peak in  $[H_2SO_4]$  and peak concentration of measurable newly formed nanoparticles (down to 3 nm). The time of this delay is consistent with  $t_0$  shown in Figure 1. Note that  $t_0$  shown above are based on IMN theory that has taken into account the enhanced growth rate of charged clusters. Without such enhancement (as in the cases of THN and BHN),  $t_0$  are expected to be larger. For relative low values of  $[H_2SO_4]$  ( $< \sim 1 \times 10^7/cm^3$ ) and when  $[H_2SO_4]$  changes rapidly,  $t_0$  may be much bigger than the time period that  $[H_2SO_4]$  can be treated as constant. Under such conditions, the assumption that pseudo-steady state concentration of critical clusters is achieved instantaneously following the

changes of  $[H_2SO_4]$ , which is a key assumption of classical nucleation theories, is no longer valid.

[16] Figure 1 also indicates that  $J_{3nm}^{IMN}$  are a factor of 2–10 smaller than  $J_{D_{crit}}^{IMN}$ . The difference is due to the scavenging of nucleated clusters by pre-existing particles before they grow to 3 nm.

### 3.2. Sensitivity of Nucleation Rate to Sulfuric Acid Concentration

[17] Since nucleation rate is very sensitive to  $t$  when  $t < t_0$ , we compare  $J_{D_{crit}}^{IMN}$  at  $t = 180$  minutes to make sure that pseudo-steady states have been reached under the conditions. Figure 2 describes the dependence of pseudo-steady state nucleation rate  $J_{D_{crit}}^{IMN}$  on  $[H_2SO_4]$  at three ionization rates ( $Q = 2, 5, 10$  ion-pairs/ $cm^3 s$ ). The nucleation rates predicted with classical BHN theory ( $J^{BHN}$ ) and THN theory at two  $NH_3$  concentrations ( $J^{THN}$ ) are also shown for comparison.  $J^{THN}$  is calculated from the parameterization given by *Napari et al.* [2002].  $J^{BHN}$  are extremely small under the conditions and have been scaled by a factor of  $10^{10}$  to be displayed in the same plot. Both THN and IMN theories can predict significant nucleation under the



**Figure 2.** The dependence of pseudo-steady state nucleation rate  $J_{D_{crit}}^{IMN}$  on  $[H_2SO_4]$  at three ionization rates ( $Q = 2, 5, 10$  ion-pairs/ $cm^3 s$ ) for (a) case 1 and (b) case 2. The nucleation rates predicted with classical BHN theory ( $J^{BHN}$ ) and THN theory at two  $NH_3$  concentrations ( $J^{THN}$ , based on parameterization given by *Napari et al.*, 2002) are also shown for comparison. Note that  $J^{BHN}$  have been scaled by a factor of  $10^{10}$ . The sensitivity of nucleation rates to  $[H_2SO_4]$  can be seen from the values of  $\alpha$  in the relation  $J \propto X^\alpha$  where  $X = [H_2SO_4]$ .

conditions considered.  $J^{\text{THN}}$  is very sensitive to  $\text{NH}_3$  concentration ( $[\text{NH}_3]$ ). A factor of 10 increase in  $[\text{NH}_3]$  enhances the nucleation rate by a factor of around  $10^7$  for case 1 and  $10^6$  for case 2. THN theory can predict very high nucleation rate (even when  $[\text{H}_2\text{SO}_4]$  is very low) as long as  $[\text{NH}_3]$  is high enough. On the contrast,  $J^{\text{IMN}}$  is limited by ionization rate and the effect of ionization rate on  $J^{\text{IMN}}$  depends on  $[\text{H}_2\text{SO}_4]$ . The relative sensitivity of  $J^{\text{IMN}}$  to  $Q$  is much smaller than that of  $J^{\text{THN}}$  to  $[\text{NH}_3]$ . A detailed discussion about the effect of  $Q$  on  $J^{\text{IMN}}$  under different atmospheric conditions can be found in Yu [2002].

[18] By assuming the functional power dependence of nucleation rate on  $[\text{H}_2\text{SO}_4]$  in the form of  $J = \text{const} \times [\text{H}_2\text{SO}_4]^\alpha = \text{const} \times X^\alpha$ , we can compare the sensitivity of  $J$  to  $[\text{H}_2\text{SO}_4]$  from the values of  $\alpha$ . For the portion of curves shown in Figure 2, the value of  $\alpha$  is 6.9 for  $J^{\text{THN}}$  and 14.0 for  $J^{\text{BHN}}$  in case 1 and is 6.0 for  $J^{\text{THN}}$  and 15.7 for  $J^{\text{THN}}$  in case 2. Based on THN theory, the presence of  $\text{NH}_3$  not only significantly increases the nucleation rates but also reduces the sensitivity of nucleation rates to  $[\text{H}_2\text{SO}_4]$  (i.e., the values of  $\alpha$ ). Nevertheless,  $J^{\text{THN}}$  is still very sensitive to  $[\text{H}_2\text{SO}_4]$  ( $\alpha \geq 6$  for the conditions shown in Figure 1) which is unlikely to explain the observed  $[\text{H}_2\text{SO}_4]^2$  dependence of  $J^{\text{obs}}$  on  $[\text{H}_2\text{SO}_4]$  (i.e.,  $\alpha = 2$ ) [Weber et al., 1996].

[19] The values of  $\alpha$  for  $J_{D_{\text{crit}}}^{\text{IMN}}$  at three different values of  $[\text{H}_2\text{SO}_4]$  are also shown in Figure 2. Under the considered conditions,  $J^{\text{IMN}}$  is negligible when  $[\text{H}_2\text{SO}_4] < \sim 3 \times 10^6/\text{cm}^3$ .  $J^{\text{IMN}}$  increase rapidly with  $[\text{H}_2\text{SO}_4]$  when  $[\text{H}_2\text{SO}_4]$  is relatively low ( $\alpha > 14$  at  $[\text{H}_2\text{SO}_4]$  of  $< \sim 3 \times 10^6/\text{cm}^3$ ) but such increase slows down when  $[\text{H}_2\text{SO}_4]$  becomes higher ( $\alpha = 2$  at  $[\text{H}_2\text{SO}_4]$  of  $\sim 9 \times 10^6/\text{cm}^3$  and  $\alpha < 0.4$  at  $[\text{H}_2\text{SO}_4]$  of  $> \sim 4.5 \times 10^7/\text{cm}^3$ ). The main physics behind such a behavior is that at low  $[\text{H}_2\text{SO}_4]$  the nucleation is limited by  $[\text{H}_2\text{SO}_4]$  while at high  $[\text{H}_2\text{SO}_4]$ , nucleation is limited by ion concentration (or ionization rate). The dramatic change of  $\alpha$  values for  $J^{\text{IMN}}$  in the range of  $[\text{H}_2\text{SO}_4]$ , where most of nucleation events in the atmosphere have been observed, is quite different from that predicted based on BHN and THN theories. The  $\alpha$  value of  $\sim 2$  when  $[\text{H}_2\text{SO}_4]$  is slightly smaller than  $1 \times 10^7/\text{cm}^3$  may somehow explain the  $[\text{H}_2\text{SO}_4]^2$  dependence of  $J^{\text{obs}}$  on  $[\text{H}_2\text{SO}_4]$  [Weber et al., 1996]. However, we should be cautious about such comparison because  $[\text{H}_2\text{SO}_4]$  changes rapidly during the observed nucleation events while we keep  $[\text{H}_2\text{SO}_4]$  constant in our simulations. Some features of  $J^{\text{obs}}$ , for example the relatively high values of  $J^{\text{obs}}$  at very low  $[\text{H}_2\text{SO}_4]$  may be explained by time delays inherent in the chemical-microphysical system under investigation [Turco et al., 1998].

#### 4. Summary

[20] We first investigate the time-needed ( $t_0$ ) to establish a pseudo-steady state production rate of certain size particles when  $[\text{H}_2\text{SO}_4]$  suddenly increases to a certain level. We find that  $t_0$  is almost inversely proportional to  $[\text{H}_2\text{SO}_4]$ . In the two marine atmospheric conditions considered in this paper, to establish a pseudo-steady state production rate of particles with the size of critical embryo, it takes  $\sim 5$  minutes and  $\sim 60$  minutes at  $[\text{H}_2\text{SO}_4]$  of  $5 \times 10^7/\text{cm}^3$  and  $5 \times 10^6/\text{cm}^3$ , respectively. Note that at a given  $[\text{H}_2\text{SO}_4]$ ,  $t_0$  is shorter at a lower temperature as  $D_{\text{crit}}$

is smaller. However, in the lower atmosphere where temperature is relatively lower, the average concentration of  $\text{H}_2\text{SO}_4$  is generally lower which will increase  $t_0$ . For relative low values of  $[\text{H}_2\text{SO}_4]$  and when  $[\text{H}_2\text{SO}_4]$  changes rapidly,  $t_0$  may be much bigger than the time period that  $[\text{H}_2\text{SO}_4]$  can be treated as constant. In such cases, the assumption that pseudo-steady state distribution of critical clusters is achieved immediately following the changes in  $[\text{H}_2\text{SO}_4]$ , on which the classical nucleation theories are based, is no longer valid.

[21] The sensitivity of nucleation rates predicted by IMN theory ( $J^{\text{IMN}}$ ) to  $[\text{H}_2\text{SO}_4]$  has been investigated and compared to that of nucleation rates predicted by BHN and THN theories. Under the conditions considered in this study,  $J^{\text{IMN}}$  is very sensitive to  $[\text{H}_2\text{SO}_4]$  when  $[\text{H}_2\text{SO}_4]$  is relatively low ( $< \sim 7 \times 10^6/\text{cm}^3$ ) but such sensitivity reduces as  $[\text{H}_2\text{SO}_4]$  increases. If we assume that nucleation rate is proportional to  $[\text{H}_2\text{SO}_4]^\alpha$ , the value of  $\alpha$  for  $J^{\text{IMN}}$  is  $> 14$ ,  $2$ ,  $< 0.4$  at  $[\text{H}_2\text{SO}_4]$  of  $3 \times 10^6$ ,  $9 \times 10^6$ ,  $4.5 \times 10^7/\text{cm}^3$ , respectively. The dramatic change of  $\alpha$  values for  $J^{\text{IMN}}$  in the range of  $[\text{H}_2\text{SO}_4]$  where most of nucleation events in the atmosphere have been observed is quite different from that predicted based on BHN ( $\alpha > 14$ ) and THN ( $\alpha = \sim 6-7$ ) theories.

[22] **Acknowledgments.** The author thanks I. Napari for providing the computer codes of their published parameterization used to calculate binary and ternary nucleation rates. This work was supported by the NSF under grant ATM 0104966.

#### References

- Ball, S. M., D. R. Hanson, F. L. Eisele, and P. H. McMurry, Laboratory studies of particle nucleation: Initial results for  $\text{H}_2\text{SO}_4$ ,  $\text{H}_2\text{O}$ , and  $\text{NH}_3$  vapors, *J. Geophys. Res.*, *104*, 23,709–23,718, 1999.
- Clarke, A. D., et al., Particle nucleation in the tropical boundary layer and its coupling to marine sulfur sources, *Science*, *282*, 89–92, 1998.
- Coffman, D. J., and D. A. Hegg, A preliminary study of the effect of ammonia on particle nucleation in the marine boundary layer, *J. Geophys. Res.*, *100*, 7147–7160, 1995.
- Hamill, P., R. P. Turco, C. S. Kiang, O. B. Toon, and R. C. Whitten, An analysis of various nucleation mechanisms for sulfate particles in the stratosphere, *J. Aerosol Sci.*, *13*, 561–585, 1982.
- Korhonen, P. M., et al., Ternary nucleation of  $\text{H}_2\text{SO}_4$ ,  $\text{NH}_3$ , and  $\text{H}_2\text{O}$  in the atmosphere, *J. Geophys. Res.*, *104*, 26,349–26,353, 1999.
- Nadykto, A. B., and F. Yu, Uptake of neutral polar vapor molecules by charged clusters/particles: Enhancement due to dipole-charge interaction, *J. Geophys. Res.*, under review, 2003.
- Napari, I., M. Noppel, H. Vehkamäki, and M. Kulmala, Parametrization of ternary nucleation rates for  $\text{H}_2\text{SO}_4$ - $\text{NH}_3$ - $\text{H}_2\text{O}$  vapors, *J. Geophys. Res.*, *107*(D19), 4381, doi:10.1029/2002JD002132, 2002.
- Raes, F., J. Augustin, and R. Vandingenen, The role of ion-induced aerosol formation in the lower atmosphere, *J. Aerosol Sci.*, *17*, 466–470, 1986.
- Reiter, R., *Phenomena in Atmospheric and Environmental Electricity*, Elsevier, New York, 1992.
- Turco, R. P., J. X. Zhao, and F. Yu, A new source of tropospheric aerosols: Ion-ion recombination, *Geophys. Res. Lett.*, *25*, 635–638, 1998.
- Weber, R. J., J. J. Marti, P. H. McMurray, F. L. Eisele, D. J. Tanner, and A. Jefferson, Measured atmospheric new particle formation rates: Implications for nucleation mechanisms, *Chem. Eng. Comm.*, *151*, 53–62, 1996.
- Yu, F., Altitude variations of cosmic ray induced production of aerosols: Implications for global cloudiness and climate, *J. Geophys. Res.*, *107*(A7), 1118, doi:10.1029/2001JA000248, 2002.
- Yu, F., and R. P. Turco, From molecular clusters to nanoparticles: The role of ambient ionization in tropospheric aerosol formation, *J. Geophys. Res.*, *106*, 4797–4814, 2001.
- Yu, F., and R. P. Turco, Ultrafine aerosol formation via ion-mediated nucleation, *Geophys. Res. Lett.*, *27*, 883–886, 2000.