

Publications, Haijun Chen

- Chen K, Zuo D, Wang S, **Chen H*** (2018). Kir2 inward rectification-controlled precise and dynamic balances between Kir2 and HCN currents initiate pacemaker activity. *FASEB Journal* (In Press).
- Xiong J, Liu X, Gong Y, Zhang P, Qiang S, Zhao Q, Guo R, Qian Y, Wang L, Zhu L, Wang R, Hao Z, Wen H, Zhang J, Tang K, Zang WF, Yuchi Z, **Chen H**, Chen SRW, Zheng W, Wang SQ, Xu YW, Liu Z* (2018) Pathogenic mechanism of a catecholaminergic polymorphic ventricular tachycardia causing-mutation in cardiac calcium release channel RyR2. *Journal of Molecular and Cellular Cardiology* (In press).
- Chen K, Zuo D, Liu Z, **Chen H*** (2017). Kir2.1 channels set two levels of resting membrane potential with inward rectification. *Pflugers Arch: European Journal of Physiology*. 2018, 470(4): 599-611.
- Zuo D, Chen K, Zhou M, Liu Z, **Chen H*** (2017). Kir2.1 and K2P1 channels reconstitute two levels of resting membrane potential in cardiomyocytes. *Journal of Physiology-London*. 2017, 595(15): 5129-5142
- Wang W, Kiyoshi CM, Du Y, Ma B, Alford CC, **Chen H** & Zhou M. (2016). mGluR3 Activation Recruits Cytoplasmic TWIK-1 Channels to Membrane that Enhances Ammonium Uptake in Hippocampal Astrocytes. *Molecular neurobiology*. 2016, 53(9):6169-6182. PubMed PMID: 26553349.
- Du Y, Kiyoshi CM, Wang Q, Wang W, Ma B, Alford CC, Zhong S, Wan Q, **Chen H**, Lloyd EE, Bryan RM, Jr., Zhou M. (2016) Genetic Deletion of TREK-1 or TWIK-1/TREK-1 Potassium Channels does not Alter the Basic Electrophysiological Properties of Mature Hippocampal Astrocytes In Situ. *Frontiers in Cellular Neuroscience*. 2016;10:13. doi: 10.3389/fncel.2016.00013. PubMed PMID: 26869883; PubMed Central PMCID: PMC4738265.
- **Chen H**, Zuo D, Zhang J, Zhou M, and Liqun Ma (2014) Classification of two-pore domain potassium channels based on rectification under quasi-physiological ionic conditions. *Channels*. In press.
- **Chen H**, Chatelain FC, Lesage F (2014) Altered and dynamic ion selectivity of K⁺ channels in cell development and excitability. *Trends in Pharmacological Sciences* 35(9): 461-469.
- Wang, W., A. Putra, G. P. Schools, B. Ma, **H. Chen**, L. K. Kaczmarek, J. Barhanin, F. Lesage, and M. Zhou (2013) The contribution of TWIK-1 channels to astrocyte K(+) current is limited by retention in intracellular compartments. *Frontiers in cellular neuroscience* 7:24.
- Barcia G, Fleming MR, Deligniere A, Gazula VR, Brown MR, Langouet M, **Chen H**, Kronengold J, Abhyankar A, Cilio R, Nitschke P, Kaminska A, Boddaert N, Casanova JL, Desguerre I, Munnich A, Dulac O, Kaczmarek LK, Colleaux L, Nabbout R (2012). De novo gain-of-function KCNT1 channel mutations cause malignant migrating partial seizures of infancy. *Nature genetics*. 2012;44(11):1255-9.
- Ma L, Zhang X, Zhou M, **Chen H** (2012) Acid-sensitive TWIK and TASK potassium channels change ion selectivity and become permeable to sodium in extracellular acidification. *Journal of Biological Chemistry* 287: 37145-37153.
- Ma L, Xie Y, Zhou M, **Chen H** (2012) Silent TWIK-1 potassium channels conduct monovalent cation currents. *Biophysical Journal* 102:L34-L36.
- Ma, L., Zhang, X. and **Chen, H.** (2011) TWIK-1 Two-Pore Domain Potassium Channels Change Ion Selectivity and Conduct Inward Leak Sodium Currents in Hypokalemia. *Science Signaling*, 4(176), ra37. [[abstract](#)] [[full text](#)] [[pdf reprint](#)][[featured in](#)]
- Gazula VR, Strumbos JG, Mei X, **Chen H**, Rahner C, Kaczmarek LK (2010). Localization of Kv1.3 channels in presynaptic terminals of brainstem auditory neurons. *Journal of Comparative*

Neurology. 518(16):3205-20.

- Zhou, M., Xu, G., Xie, M., Zhang, X., Schools, G.P., Ma, L., Kimelberg, H.K., and **Chen, H.** (2009) TWIK-1 and TREK-1 are potassium channels contributing significantly to astrocyte passive conductance in rat hippocampal slices. *Journal of Neuroscience* 29(26): 8551-8564
- **Chen, H.**, Kronengold, J., Yan, J., Gazula, V.R., Ferreira, G., Yang, Y., Bhattacharjee, A., Sigworth, F.J., Salkoff, L., and Kaczmarek, L.K. (2009) The N-terminal domain of Slack determines the formation and functional expression of Slick/Slack heteromeric sodium-activated potassium channels. *Journal of Neuroscience*. 29(17): 5654-5665
- **Chen H.** and Goldstein S. A. (2007) Serial perturbation of MinK in IKs implies an α -helical transmembrane span traversing the channel corpus. *Biophysical Journal*, 93(7): 2332-40.
- **Chen H.**, C. von Hehn, L.K. Kaczmarek, L. R. Ment, B.R. Pober, and F.M. Hisama. (2007) Functional Analysis of a Novel Potassium Channel (KCNA1) Mutation in Hereditary Myokymia. *Neurogenetics* 8(2):131-5.
- Sun, A.-Q., Balasubramanian N., **Chen H.**, M. Shahid M., Suchy FJ. (2006). Identification of functionally relevant residues of the rat ileal apical sodium dependent bile acid cotransporter (rASBT). *Journal of Biological Chemistry* 281, 16410-8.
- Barbier, J., Lamthanh, H., Le Gall F., Favreau P., Benoit E., **Chen, H.**, Gilles N., Ilan N., Heinemann, S.H., Gurevitz M., Gordon D., Ménez A., and Molgó, J. (2004). A δ -conotoxin from *Conus ermineus* venom inhibits inactivation in vertebrate neuronal Na⁺ channels, but not in skeletal and cardiac muscles. *Journal of Biological Chemistry* 279, 4680-4685.
- **Chen H.** Kim, L.A., Rajan S., Xu S., and Goldstein, S. A. (2003). Charybdotoxin binding in the IKs pore demonstrates two MinK subunits in each channel complex. *Neuron* 40, 15-23.
- **Chen, H.**, Sesti, F., and Goldstein, S. A. (2003). Pore- and State-Dependent Cadmium Block of IKs Channels Formed with MinK-55C and Wild-Type KCNQ1 Subunits. *Biophysical Journal* 84, 3679-3689.
- **Chen, H.**, Lu, S., Leipold, E., Gordon, D., Hansel, A., and Heinemann, S. H. (2002). Differential sensitivity of sodium channels from the central and peripheral nervous system to the scorpion toxins Lqh-2 and Lqh-3. *European Journal of Neuroscience* 16, 767-770.
- **Chen, H.**, and Heinemann, S. H. (2001). Interaction of scorpion alpha-toxins with cardiac sodium channels: binding properties and enhancement of slow inactivation. *Journal of General Physiology* 117, 505-518.
- Gilles, N., Leipold, E., **Chen, H.**, Heinemann, S. H., and Gordon, D. (2001). Effect of depolarization on binding kinetics of scorpion alpha-toxin highlights conformational changes of rat brain sodium channels. *Biochemistry* 40, 14576-14584.
- **Chen, H.**, Gordon, D., and Heinemann, S. H. (2000). Modulation of cloned skeletal muscle sodium channels by the scorpion toxins Lqh II, Lqh III, and Lqh alphaIT. *Pflugers Arch: European Journal of Physiology* 439, 423-432.
- Gilles, N., **Chen, H.**, Wilson, H., Le Gall, F., Montoya, G., Molgo, J., Schonherr, R., Nicholson, G., Heinemann, S. H., and Gordon, D. (2000). Scorpion alpha and alpha-like toxins differentially interact with sodium channels in mammalian CNS and periphery. *European Journal of Neuroscience* 12, 2823-2832.
- **Chen, H.**, Kubo, Y., Hoshi, T., and Heinemann, S.H. (1998). Cyclosporin A selectively reduces functional expression of Kir2.1 potassium channels in *Xenopus* oocytes. *FEBS letters* 422, 307-310.